

Singular Vectors of the $N = 2$ Superconformal Algebra

Matthias Dörrzapf *

*Department of Applied Mathematics and Theoretical Physics
University of Cambridge, Silver Street
Cambridge, CB3 9EW, U. K.***ABSTRACT**

We derive a set of recursion formulae to construct singular vectors for the $N = 2$ (untwisted) algebra, by using the approach of Bauer, di Francesco, Itzykson and Zuber. Applying these formulae, we obtain explicit expressions for the charged singular vectors and for a class of uncharged singular vectors.

March 1994

*e-mail: M.Doerrzapf@amtp.cam.ac.uk

1. Introduction

Recent developments in string theory have shown that the $N = 2$ critical string provides a consistent quantum theory of self-dual gravity in four dimensions^{1,2}, and furthermore the infinite spectrum of particles which appears in other string theories turns out to be a finite spectrum for $N = 2$ critical strings. The interest in $N = 2$ superstring theory is also motivated by the conjecture of Atiyah and Ward³, that all the bosonic integrable systems in lower dimensions can be generated by self-dual Yang-Mills theory in four dimensional space-time with signature $(2, 2)$. This leads to the suggestion that the $N = 2$ critical string should give us insight in integrable systems. Furthermore, the $N = 2$ superstring seems to be closely related to the theory of two dimensional black holes⁴. A treatment of canonical quantum supergravity in three dimensions and the special rôle of the $N = 2$ superstring can be found, for instance, in ref. 5.

It is not only the importance of $N = 2$ critical superstring theory that makes $N = 2$ superconformal theories particularly interesting; there are also applications in two dimensional critical phenomena. Starting with the paper of Belavin, Polyakov and Zamolodchikov⁶, many statistical models at their critical points have been identified as conformally invariant theories⁷⁻⁹. Furthermore, Friedan, Qiu and Shenker were able to identify the tricritical Ising model^a as a $N = 1$ superconformal model¹¹. This raised the interesting question if $N = 2$ superconformal theories could be found to describe some statistical models at their critical points, and indeed it was shown in ref. 12 that under certain circumstances $O(2)$ Gaussian models are $N = 2$ superconformally invariant.

The physically important representations of the $N = 2$ superconformal algebra are the irreducible highest weight representations. In order to understand their structure, it is helpful to find explicit expressions for the singular vectors appearing in the Verma modules. Furthermore, since these singular vectors decouple from the theory, they imply differential equations for the correlation functions. For the Virasoro algebra explicit formulae for some singular vectors were first given by Benoit and Saint-Aubin¹³ and later for all singular vectors by Bauer et al.^{14,15} and by Kent¹⁶. There are basically three different methods known: the “*fusion method*” used by Bauer et al.^{14,15}, the “*analytic continuation method*” developed by Malikov, Feigin and Fuchs for Kac-Moody algebras¹⁷ and extended by Kent to the Virasoro algebra¹⁶ and the method of Ganchev and Petkova which uses the Knizhnik-Zamolodchikov equation to transform a Kac-Moody singular vector into a Virasoro one¹⁸. The fusion method has been used for various other chiral algebras as well: for the $N = 1$ Neveu-Schwarz algebra by Benoit and Saint-Aubin¹⁹ and Huang et al.²⁰, for the $N = 1$ Ramond algebra by Watts²¹, for the W_3 algebra by Bowcock and Watts²², for the affine algebra $A_1^{(1)}$ by Bauer and Sochen²³ and for the WBC_2 algebra by Bajnok²⁴. Recently the analytic continuation method was used by Bajnok²⁵ to find WA_2 singular vectors.

We shall show in this paper that the fusion method can be used successfully in the $N = 2$ case. In section 2 we will introduce the notation and derive the $N = 2$ descent equations, arising from the operator product expansion of $N = 2$ superconformal fields. In the first part

^aExperimentally realised in a ⁴He monolayer absorbed on Kr-plated graphite¹⁰.

of section 3 we use heuristic arguments to derive a set of recursion formulae which allow us to compute singular vectors at high levels using known singular vectors at lower levels. A rigorous proof for these recursion formulae is given in the second part of section 3. In the fourth section we use the recursion formulae to find explicit expressions for all charged singular vectors and for a class of uncharged singular vectors which are the analogues of the Benoit and Saint-Aubin singular vectors in the Virasoro case. Finally we investigate the dependence of the recursion on the position of the fusion point.

2. $N = 2$ Superconformal Theories

2.1. Notation

The algebra of chiral superconformal transformations in $N = 2$ superconformal space²⁶ is generated by the super stress-energy tensor $T(Z_1)$, where Z_1 denotes a superpoint $(z_1, \theta_{1,1}, \theta_{1,2})$. Superconformal invariance of the theory determines the operator product expansion (OPE) or short distance expansion of $T(Z)$ with itself:

$$T(Z_1) T(Z_2) = -\frac{c}{12Z_{12}^2} + \left(-\frac{\theta_{12,1}\theta_{12,2}}{Z_{12}^2} + \frac{1}{2} \frac{\theta_{12,2}D_{2,1} - \theta_{12,1}D_{2,2}}{Z_{12}} - \frac{\theta_{12,1}\theta_{12,2}}{Z_{12}} \partial_{z_2} \right) T(Z_2), \quad (2.1)$$

where we have used $\theta_{12,i} = \theta_{1,i} - \theta_{2,i}$ and $Z_{12} = z_1 - z_2 - \theta_{1,i}\theta_{2,i}$ as superintegrals of unity and $D_{i,j} = \frac{\partial}{\partial \theta_{i,j}} + \theta_{i,j} \frac{\partial}{\partial z}$ as superderivatives. If we do not need to label the superpoint, we shall just write θ_i and D_i .

Expanding $T(Z)$ in its modes allows us to find the symmetry algebra generators^b:

$$\begin{aligned} T(Z_1) &= -\theta_{12,1}\theta_{12,2} \sum_{m \in \mathbb{Z}} Z_{12}^{-m-2} L_m(Z_2) - \frac{1}{2} \sum_{r \in \mathbb{Z} + \frac{1}{2}} Z_{12}^{-r-\frac{3}{2}} \theta_{12,2} G_r^1(Z_2) \\ &\quad + \frac{1}{2} \sum_{r \in \mathbb{Z} + \frac{1}{2}} Z_{12}^{-r-\frac{3}{2}} \theta_{12,1} G_r^2(Z_2) - \frac{1}{2} i \sum_{m \in \mathbb{Z}} T_m Z_{12}^{-m-1}(Z_2) \end{aligned} \quad (2.2)$$

$$\begin{aligned} L_m(Z_2) &= \oint \frac{dZ_1}{2\pi i} Z_{12}^{m+1} T(Z_1) \\ G_r^i(Z_2) &= 2 \oint \frac{dZ_1}{2\pi i} \theta_{12,i} Z_{12}^{r+\frac{1}{2}} T(Z_1) \\ T_m(Z_2) &= -2i \oint \frac{dZ_1}{2\pi i} \theta_{12,1}\theta_{12,2} Z_{12}^m T(Z_1). \end{aligned} \quad (2.3)$$

If we use complex co-ordinates for the odd generators

$$\theta^\pm = \frac{1}{\sqrt{2}}(\theta_1 \pm i\theta_2) \quad (2.4)$$

^bWe put $Z_2 = 0$ to obtain the field-state correspondence at the origin and we use L_m for $L_m(0)$ etc.

$$G^\pm = \frac{1}{\sqrt{2}}(G^1 \pm iG^2) \quad (2.5)$$

$$D^\pm = \frac{1}{\sqrt{2}}(D_1 \pm iD_2) = \frac{\partial}{\partial\theta^\mp} + \theta^\pm \frac{\partial}{\partial z}, \quad (2.6)$$

we find for the symmetry algebra $\mathcal{A}$ a decomposition in two odd fields, the Virasoro algebra and a $U(1)$ Kac-Moody algebra with the commutation relations:

$$\begin{aligned} [L_m, L_n] &= (m-n)L_{m+n} + \frac{c}{12}(m^3-m)\delta_{m+n,0} \\ [L_m, G_r^\pm] &= \left(\frac{1}{2}m-r\right)G_{m+r}^\pm \\ [L_m, T_n] &= -nT_{m+n} \\ [T_m, T_n] &= \frac{1}{3}cm\delta_{m+n,0} \\ [T_m, G_r^\pm] &= \pm G_{m+r}^\pm \\ \{G_r^+, G_s^-\} &= 2L_{r+s} + (r-s)T_{r+s} + \frac{c}{3}\left(r^2 - \frac{1}{4}\right)\delta_{r+s,0} \\ \{G_r^+, G_s^+\} &= \{G_r^-, G_s^-\} = 0, \quad m, n \in \mathbb{Z}; \quad r, s \in \mathbb{Z} + \frac{1}{2}. \end{aligned} \quad (2.7)$$

The primary fields $\Phi(Z)$ in the theory are those transforming homogeneously under superconformal transformations. This leads to a short distance expansion of $T(Z)$ and $\Phi(Z)$ of the form:

$$\begin{aligned} T(Z_1)\Phi(Z_2) &= -\frac{h\theta_{12,1}\theta_{12,2}}{Z_{12}^2}\Phi(Z_2) + \frac{1}{2}\frac{\theta_{12,2}D_{2,1} - \theta_{12,1}D_{2,2}}{Z_{12}}\Phi(Z_2) \\ &\quad - \frac{\theta_{12,1}\theta_{12,2}}{Z_{12}}\partial_{z_2}\Phi(Z_2) - \frac{q}{2Z_{12}}i\Phi(Z_2). \end{aligned} \quad (2.8)$$

We call h the *conformal weight* of Φ and q its *conformal charge*, corresponding to the scaling dimensions of L_0 and T_0 transformations respectively. Using contour integral methods we obtain the infinitesimal transformations for all generators:

$$\begin{aligned} [L_m, \Phi(Z)] &= \left\{ h(m+1)z^m + \frac{1}{2}(m+1)z^m(\theta^+D^- + \theta^-D^+) + z^{m+1}\partial_z \right. \\ &\quad \left. + \frac{q}{2}\theta^+\theta^-z^{m-1}m(m+1) \right\} \Phi(Z) \\ [G_r^\pm, \Phi(Z)] &= \left\{ 2h\left(r + \frac{1}{2}\right)\theta^\pm z^{r-\frac{1}{2}} - z^{r+\frac{1}{2}}D^\pm \pm \theta^+\theta^-\left(r + \frac{1}{2}\right)z^{r-\frac{1}{2}}D^\pm \right. \\ &\quad \left. + 2\theta^\pm z^{r+\frac{1}{2}}\partial_z \pm q\theta^\pm z^{r-\frac{1}{2}}\left(r + \frac{1}{2}\right) \right\} \Phi(Z) \\ [T_m, \Phi(Z)] &= \left\{ 2h\theta^+\theta^-mz^{m-1} + z^m(\theta^-D^+ - \theta^+D^-) + 2\theta^+\theta^-z^m\partial_z \right. \\ &\quad \left. + qz^m \right\} \Phi(Z). \end{aligned} \quad (2.9)$$

We should note at this stage that the $N = 2$ superconformal algebra we consider is known as the $N = 2$ Neveu-Schwarz or antiperiodic algebra. In ref. 27 it has been shown that it is isomorphic to the $N = 2$ Ramond or periodic algebra which makes a separate discussion

redundant. The $N = 2$ twisted algebra does not fit in this framework since the corresponding fields are not $N = 2$ superconformal in our sense, however a discussion of its singular vectors was given by Semikhatov²⁸. A definition of $N = 2$ superconformal theories can be found for instance in refs. 29, 30.

$\mathcal{A}$ can be decomposed in the usual way: $\mathcal{A} = \mathcal{A}_- \oplus \mathcal{H} \oplus \mathcal{A}_+$, where $\mathcal{H} = \text{span}\{L_0, T_0\}$ is the Cartan subalgebra, and^c $\mathcal{A}_\pm = \text{span}\{L_{\pm n}, T_{\pm n}, G_{\pm r}^+, G_{\pm r}^- : n \in \mathbb{N}, r \in \mathbb{N}_0 + \frac{1}{2}\}$. A highest weight vector $|h, q\rangle$ is a simultaneous eigenvector of $\mathcal{H}$ with L_0 and T_0 eigenvalues h and q respectively, and $\mathcal{A}_+ |h, q\rangle = 0$. It is easy to see that a primary field $\Phi_{h,q}$ generates a highest weight vector $|h, q\rangle$ on the vacuum: $|h, q\rangle = \Phi_{h,q}(0) |0\rangle$. If $U(\mathcal{A})$ denotes the universal enveloping algebra of $\mathcal{A}$ then the Verma module $\mathcal{V}_{h,q}$ is defined as $\mathcal{V}_{h,q} = U(\mathcal{A}) \otimes_{\mathcal{H} \oplus \mathcal{A}_+} |h, q\rangle$. This is the representation of $\mathcal{A}$ with the basis

$$\left\{ L_{-i_I} \dots L_{-i_1} G_{-j_+}^+ \dots G_{-j_1}^+ G_{-j_-}^- \dots G_{-j_1}^- T_{-k_K} \dots T_{-k_1} |h, q\rangle ; \right. \\ \left. i_I \geq \dots \geq i_1 \geq 1, j_{J^+}^+ > \dots > j_1^+ \geq \frac{1}{2}, j_{J^-}^- > \dots > j_1^- \geq \frac{1}{2}, k_K \geq \dots \geq k_1 \geq 1 \right\}.$$

Vectors in a Verma module which are not multiples of the highest weight vector but satisfy the highest weight vector conditions are called *singular vectors*, i.e. $\Psi_{\tilde{h}, \tilde{q}} \in \mathcal{V}_{h,q}$ is called singular if $L_0 \Psi_{\tilde{h}, \tilde{q}} = \tilde{h} \Psi_{\tilde{h}, \tilde{q}}$, $T_0 \Psi_{\tilde{h}, \tilde{q}} = \tilde{q} \Psi_{\tilde{h}, \tilde{q}}$ and $\mathcal{A}_+ \Psi_{\tilde{h}, \tilde{q}} = 0$.

2.2. Commutation of fields and algebra elements

In this subsection we shall define a representation of the $N = 2$ algebra which allows us to interchange fields and algebra elements:

$$\Phi_{h,q}(Z)X = \sigma(X)\Phi_{h,q}(Z) \quad , \quad X \in \mathcal{A}.$$

Using the equations (2.9) we find:

$$\begin{aligned} \sigma(L_m) &= L_m - h(m+1)z^m - z^{m+1}\partial_z - \frac{1}{2}(m+1)z^m(\theta^+ D^- + \theta^- D^+) \\ &\quad - \frac{q}{2}m(m+1)\theta^+ \theta^- z^{m-1} \\ \sigma(G_r^\pm) &= G_r^\pm - 2h(r + \frac{1}{2})\theta^\pm z^{r-\frac{1}{2}} \mp (r + \frac{1}{2})\theta^+ \theta^- z^{r-\frac{1}{2}} D^\pm \\ &\quad + z^{r+\frac{1}{2}} D^\pm - 2\theta^\pm z^{r+\frac{1}{2}} \partial_z \mp q(r + \frac{1}{2})z^{r-\frac{1}{2}} \theta^\pm \\ \sigma(T_m) &= T_m - 2hm\theta^+ \theta^- z^{m-1} - z^m(\theta^- D^+ - \theta^+ D^-) - 2\theta^+ \theta^- z^m \partial_z - qz^m. \end{aligned} \tag{2.10}$$

Trivially, this defines a representation on primary fields. However, it can be shown by direct computation that it defines a representation in general.

It is convenient to choose a basis l_m, g_r^i and t_m for the algebra $\mathcal{A}$ such that σ does not involve any derivatives on the basis. In the manner of Friedan et al.¹¹ we find that σ terms

^cWe write $\mathbb{N}$ for $\{1, 2, 3, \dots\}$ and $\mathbb{N}_0$ for $\{0, 1, 2, \dots\}$.

involving derivatives arise from singular contributions to the OPE. The only singular terms we should allow are of the form $\frac{\theta_{12,1}\theta_{12,2}}{Z_{12}}$. A possible choice is:

$$\begin{aligned} l_m(Z_3) &= \oint_{\mathcal{C}} \frac{dZ_1}{2\pi i} Z_{12}^m Z_{13} T(Z_1) \\ g_r^i(Z_3) &= 2 \oint_{\mathcal{C}} \frac{dZ_1}{2\pi i} Z_{12}^{r-\frac{1}{2}} Z_{13} T(Z_1) \\ t_m(Z_3) &= -2i \oint_{\mathcal{C}} \frac{dZ_1}{2\pi i} Z_{12}^{m-1} Z_{13} T(Z_1), \end{aligned} \quad (2.11)$$

where $\mathcal{C}$ is a super contour about the super points Z_2 and Z_3 , and we shall set $Z_3 = 0$.

For $m \in \mathbb{N}$ and $r \in \mathbb{N}_0 + \frac{1}{2}$ we shall use^d:

$$\begin{aligned} l_m &= L_m - zL_{m-1} + \frac{1}{2}(\theta^- G_{m-\frac{1}{2}}^+ + \theta^+ G_{m-\frac{1}{2}}^-) & \sigma(l_m) &= l_m - hz^m \\ g_r^\pm &= G_r^\pm - zG_{r-1}^\pm \mp \theta^\pm T_{r-\frac{1}{2}} & \sigma(g_r^\pm) &= g_r^\pm - 2h\theta^\pm z^{r-\frac{1}{2}} \\ t_m &= T_m - zT_{m-1} & \sigma(t_m) &= t_m - 2h\theta^+\theta^- z^{m-1}, \end{aligned} \quad (2.12)$$

whilst for $-m \in \mathbb{N}$ and $-r \in \mathbb{N}_0 + \frac{1}{2}$ we use:

$$\begin{aligned} l_m &= -L_m + \frac{1}{z}L_{m+1} + \frac{1}{2z}(\theta^- G_{m+\frac{1}{2}}^+ + \theta^+ G_{m+\frac{1}{2}}^-) & \sigma(l_m) &= l_m - hz^m \\ g_r^\pm &= -G_r^\pm + \frac{1}{z}G_{r+1}^\pm \mp \frac{1}{z}\theta^\pm T_{r+\frac{1}{2}} & \sigma(g_r^\pm) &= g_r^\pm - 2h\theta^\pm z^{r-\frac{1}{2}} \\ t_m &= -T_m + \frac{1}{z}T_{m+1} & \sigma(t_m) &= t_m - 2h\theta^+\theta^- z^{m-1}, \end{aligned} \quad (2.13)$$

and finally we have to add L_0 and T_0 to obtain a basis.

2.3. $N = 2$ descent equations

Given an OPE of two primary fields we can derive necessary conditions for the action of the algebra on the expansion coefficients in the spirit of Belavin, Polyakov and Zamolodchikov⁶ who derived the so called *descent equations* for the Virasoro case. We start from the most general expression for the OPE multiplied by the vacuum $|0\rangle$

$$\begin{aligned} \Phi_{h_1, q_1}(Z) \Phi_{h_2, q_2}(0) |0\rangle &= \sum_{j \in \mathcal{J}} C_{12}^j z^{h_j - h_1 - h_2} |\psi_j(Z)\rangle \\ &+ \theta^+ \sum_{j \in \mathcal{J}} C_{12}^{j-} z^{h_j - h_1 - h_2 - \frac{1}{2}} |\psi_j^-(Z)\rangle \\ &+ \theta^- \sum_{j \in \mathcal{J}} C_{12}^{j+} z^{h_j - h_1 - h_2 - \frac{1}{2}} |\psi_j^+(Z)\rangle \\ &+ \theta^+ \theta^- \sum_{j \in \mathcal{J}} \bar{C}_{12}^j z^{h_j - h_1 - h_2 - 1} |\bar{\psi}_j(Z)\rangle, \end{aligned} \quad (2.14)$$

^dIt is worth remarking that the G 's and θ 's anticommute.

where j runs through all the conformal families.

The conformal families $|\psi_j(Z)\rangle$, $|\psi_j^-(Z)\rangle$, $|\psi_j^+(Z)\rangle$ and $|\overline{\psi}_j(Z)\rangle$ can be written as power series in z

$$\begin{aligned} |\psi_j(Z)\rangle &= \sum_{n \in \mathbb{N}_0} z^n |h_j + n, q_j\rangle + \theta^+ \sum_{r \in \mathbb{N}_0 + \frac{1}{2}} z^{r - \frac{1}{2}} |h_j + r, q_j - 1\rangle \\ &\quad + \theta^- \sum_{r \in \mathbb{N}_0 + \frac{1}{2}} z^{r - \frac{1}{2}} |h_j + r, q_j + 1\rangle + \theta^+ \theta^- \sum_{n \in \mathbb{N}_0} z^{n-1} \overline{|h_j + n, q_j\rangle}, \end{aligned} \quad (2.15)$$

where $|h_j, q_j\rangle$ denotes the highest weight vector and $|h_j + n, q_j(\pm 1)\rangle$ its descendants; similarly for $|\psi_j^-(Z)\rangle$, $|\psi_j^+(Z)\rangle$ and $|\overline{\psi}_j(Z)\rangle$.

We apply the basis elements of $\mathcal{A}$ to equation (2.14), use the commutation relations (2.9) and take in account that $\Phi_{h_2, q_2}(0) |0\rangle$ is a highest weight state. After performing the differentiations and comparing the coefficients we find necessary conditions for the expansion coefficients in (2.15). We notice that the coefficients C_{12}^{j-} , C_{12}^{j+} and $\overline{C}_{12}^j$ have to vanish necessarily, whilst the C_{12}^j can be non trivial if $q_j = q_1 + q_2$. In this case we find the descent equations to be:

$$\begin{aligned} L_m |h_j + n + m, q_j\rangle &= (mh_1 + h_j - h_2 + n) |h_j + n, q_j\rangle \\ L_m |h_j + r + m, q_j \pm 1\rangle &= (h_1 m + h_j - h_2 + \frac{m}{2} + r) |h_j + r, q_j \pm 1\rangle \\ L_m \overline{|h_j + n + m, q_j\rangle} &= (h_1 m + h_j - h_2 + m + n) \overline{|h_j + n, q_j\rangle} \\ &\quad + \frac{q_1}{2} m(m+1) |h_j + n, q_j\rangle \\ G_s^+ |h_j + r + s, q_j\rangle &= - |h_j + r, q_j + 1\rangle \\ G_s^+ |h_j + n + s, q_j - 1\rangle &= -(2h_1 s + h_j - h_2 + n + q_1(s + \frac{1}{2})) |h_j + n, q_j\rangle \\ &\quad - \overline{|h_j + n, q_j\rangle} \\ G_s^+ |h_j + n + s, q_j + 1\rangle &= 0 \\ G_s^+ \overline{|h_j + r + s, q_j\rangle} &= (2h_1 s + h_j - h_2 + r + s + q_1(s + \frac{1}{2})) |h_j + r, q_j + 1\rangle \\ G_s^- |h_j + r + s, q_j\rangle &= - |h_j + r, q_j - 1\rangle \\ G_s^- |h_j + n + s, q_j - 1\rangle &= 0 \\ G_s^- |h_j + n + s, q_j + 1\rangle &= -(2h_1 s + h_j - h_2 + n - q_1(s + \frac{1}{2})) |h_j + n, q_j\rangle \\ &\quad + \overline{|h_j + n, q_j\rangle} \\ G_s^- \overline{|h_j + r + s, q_j\rangle} &= -(2h_1 s + h_j - h_2 + r + s - q_1(s + \frac{1}{2})) |h_j + r, q_j - 1\rangle \\ T_m |h_j + n + m, q_j\rangle &= q_1 |h_j + n, q_j\rangle \\ T_m |h_j + r + m, q_j \pm 1\rangle &= (\pm 1 + q_1) |h_j + r, q_j \pm 1\rangle \\ T_m \overline{|h_j + n + m, q_j\rangle} &= 2h_1 m |h_j + n, q_j\rangle + q_1 \overline{|h_j + n, q_j\rangle} \\ L_0 |h_j + n, q_j\rangle &= (h_j + n) |h_j + n, q_j\rangle \\ L_0 |h_j + r, q_j \pm 1\rangle &= (h_j + r) |h_j + r, q_j \pm 1\rangle \\ L_0 \overline{|h_j + n, q_j\rangle} &= (h_j + n) \overline{|h_j + n, q_j\rangle} \\ T_0 |h_j + n, q_j\rangle &= q_j |h_j + n, q_j\rangle \\ T_0 |h_j + r, q_j \pm 1\rangle &= (q_j \pm 1) |h_j + r, q_j \pm 1\rangle \\ T_0 \overline{|h_j + n, q_j\rangle} &= q_j \overline{|h_j + n, q_j\rangle}, \end{aligned} \quad (2.16)$$

where we have put $q_j = q_1 + q_2$ and $m \in \mathbb{N}$, $n \in \mathbb{N}_0$, $r, s \in \mathbb{N}_0 + \frac{1}{2}$.

In theory, we only have to solve this system of infinitely many equations for infinitely many unknowns to find out which conformal families contribute to the right hand side of equation (2.14) and this would determine the OPE completely. However, this seems impractical and we have to rely on other methods to determine the *fusion rules*^{29–31}.

2.4. The $N = 2$ determinant formula

The determinant formula for the $N = 2$ superconformal (antiperiodic) algebra was first given by Boucher, Friedan and Kent³². If we use the parametrisation:

$$\begin{aligned} c &= 3 - 3t \\ h_{r,s} &= \frac{r^2 - 1}{8}t - \frac{rs}{4} + \frac{s^2 - 1}{8t} - \frac{4q^2 - 1}{8t} \\ h_k &= kq + \frac{1}{2}t(k^2 - \frac{1}{4}), \end{aligned} \quad (2.17)$$

where r is chosen to be $\in \mathbb{Z}$, $s \in 2\mathbb{Z}$ and $k \in \mathbb{Z} + \frac{1}{2}$, then the determinant formula can be written as

$$\det(M_{n,m}(c, h, q)) \propto \prod_{\substack{1 \leq rs \leq 2n \\ r \in \mathbb{N}, s \in 2\mathbb{N}}} (h - h_{r,s})^{p(n - \frac{rs}{2}, m)} \prod_{k \in \mathbb{Z} + \frac{1}{2}} (h - h_k)^{\tilde{p}(n - |k|, m - \text{sgn}(k), k)}, \quad (2.18)$$

where $p(n - \frac{rs}{2}, m)$ and $\tilde{p}(n - |k|, m - \text{sgn}(k), k)$ are the corresponding partition functions. This makes it apparent that for every t and q , every positive integer r and every positive even s the Verma module $\mathcal{V}_{c(t), h_{r,s}(t,q), q}$ has an uncharged singular vector at level $\frac{rs}{2}$. Similarly the Verma module $\mathcal{V}_{c(t), h_k(t,q), q}$ has a singular vector at level $|k|$ with charge $\text{sgn}(k)$. By direct computation we can find for example the singular vector of the Verma module $\mathcal{V}_{c(t), h_{1,2}(t,q), q}$:

$$\Psi_{1,2} = \left(tL_{-1} + \frac{t}{q-1}G_{-\frac{1}{2}}^+ G_{-\frac{1}{2}}^- + (q+1)T_{-1} \right) |h_{1,2}, q\rangle. \quad (2.19)$$

We will use this singular vector in section 4 to compute singular vectors at higher levels recursively.

3. Recursion formulae for singular vectors

3.1. Recursion formulae

The following arguments should be understood on a heuristic level only, in order to derive the recursion formulae. In the next section we will prove these recursion formulae rigorously without using any of the heuristic arguments.

Let us consider a three point function of primary fields

$$\langle h_j, q_j | \Phi_{h_1, q_1}(Z) \Phi_{h_2, q_2}(0) | 0 \rangle, \quad (3.1)$$

where $|h_j, q_j\rangle$ is a highest weight vector generated by a primary field Φ_{h_j, q_j} . (3.1) will only be non trivial if and only if the descent equations (2.16) allow a non trivial contribution of the highest weight state $|h_j, q_j\rangle$ to the right hand side of the OPE (2.14). In this case we say we have an allowed (*even*) fusion of Φ_{h_1, q_1} and Φ_{h_2, q_2} to Φ_{h_j, q_j} . The information which three point functions are non trivial is therefore summarised in so called *fusion rules*. For the $N = 2$ case a discussion of them^e can be found in refs. 31, 29, 30. Let us furthermore assume (3.1) is non trivial which means we pick an allowed fusion and suppose we know a singular vector in the Verma module with highest weight state $\Phi_{h_2, q_2}(0)|0\rangle$, Ψ say, at level N :

$$\Psi = \mathcal{N}\Phi_{h_2, q_2}(0)|0\rangle, \quad (3.2)$$

where $\mathcal{N}$ is a linear combination of algebra generators $\in \mathcal{A}_-$.

Since the singular vectors decouple, the function $\langle h_j, q_j | \Phi_{h_1, q_1}(Z) \mathcal{N} \Phi_{h_2, q_2}(0) | 0 \rangle$ has to vanish identically^f. We use at this stage the representation σ defined in section 2.2. in order to commute $\mathcal{N}$ and $\Phi_{h_1, q_1}(Z)$:

$$0 = \langle h_j, q_j | \Phi_{h_1, q_1}(Z) \mathcal{N} \Phi_{h_2, q_2}(0) | 0 \rangle = \langle h_j, q_j | \sigma(\mathcal{N}) \Phi_{h_1, q_1}(Z) \Phi_{h_2, q_2}(0) | 0 \rangle. \quad (3.3)$$

We write $\mathcal{N}$ in the basis (2.13) before applying σ to it. This avoids any derivatives^g in $\sigma(\mathcal{N})$.

Let Π_j be the projection operator onto the conformal family generated by the primary field Φ_{h_j, q_j} . Then define the following OPEs:

$$\begin{aligned} \mathcal{F}(Z) &= \Pi_j(\Phi_{h_1, q_1}(Z) \Phi_{h_2, q_2}(0) | 0 \rangle) = \sum_{n \geq 0} z^{h_j - h_1 - h_2 + n} f_n + \theta^+ \sum_{r \in \mathbb{N}_0 + \frac{1}{2}} z^{h_j - h_1 - h_2 + r - \frac{1}{2}} f_r^- \\ &\quad + \theta^- \sum_{r \in \mathbb{N}_0 + \frac{1}{2}} z^{h_j - h_1 - h_2 + r - \frac{1}{2}} f_r^+ + \theta^+ \theta^- \sum_{n \geq 0} z^{h_j - h_1 - h_2 + n - 1} \bar{f}_n \end{aligned} \quad (3.4)$$

$$\begin{aligned} \mathcal{J}(Z) &= \sigma(\mathcal{N}) \mathcal{F}(Z) = \sum_{n \geq 0} z^{h_j - h_1 - h_2 + n - N} j_n + \theta^+ \sum_{r \in \mathbb{N}_0 + \frac{1}{2}} z^{h_j - h_1 - h_2 + r - \frac{1}{2} - N} j_r^- \\ &\quad + \theta^- \sum_{r \in \mathbb{N}_0 + \frac{1}{2}} z^{h_j - h_1 - h_2 + r - \frac{1}{2} - N} j_r^+ + \theta^+ \theta^- \sum_{n \geq 0} z^{h_j - h_1 - h_2 + n - 1 - N} \bar{j}_n \end{aligned} \quad (3.5)$$

Since we are considering an allowed fusion, $\mathcal{F}(Z)$ is non trivial. However, according to (3.3) we expect $\mathcal{J}(Z)$ to decouple. This means that it lies completely in the submodule which we have to quotient out of the Verma module $\mathcal{V}_{h_j, q_j}$ in order to get the corresponding irreducible representation. The expansion coefficients of $\mathcal{J}(Z)$ must lie in the submodule generated by a singular vector. Hence, at the level where we expect the first singular vector in $\mathcal{V}_{h_j, q_j}$ the coefficients of $\mathcal{J}(Z)$ have to vanish or give us an expression for the singular vector. In the

^eAs far as we know, the descent equations decouple for even and odd contributions and we can consider “even” and “odd” fusion separately. We will not need to use odd fusion and we write fusion for even fusion below.

^fRecall that Ψ is one representative of the zero vector in the irreducible representation with highest weight state $|h_2, q_2\rangle$.

^gHowever, we have to deal with the superderivatives now on an algebraic level in form of the descent equations.

following we derive recursion formulae for the coefficients of $\mathcal{F}(Z)$ in order to set consistently the coefficients of $\mathcal{J}(Z)$ equal to 0. We will see that in certain cases this recursion procedure breaks down at the level where we expect the first singular vector in $\mathcal{V}_{h_j, q_j}$ and the information achieved from the recursion for $\mathcal{F}(Z)$ will allow us to compute the coefficients of $\mathcal{J}(Z)$ for this level and hence the singular vector.

We write $\sigma(\mathcal{N})$ again in the standard basis:

$$\begin{aligned}
\sigma(\mathcal{N}) = & \sum_{k=0}^N z^{-N+k} S_k \\
& + \sum_{i=0}^{N-1} \sum_{k=0}^N z^{-N+k-i-1} (p_k^{1,i} L_1 + p_k^{2,i} G_{\frac{1}{2}}^+ G_{\frac{1}{2}}^- + p_{k-\frac{1}{2}}^{3-,i} G_{\frac{1}{2}}^+ + p_{k-\frac{1}{2}}^{4+,i} G_{\frac{1}{2}}^-) L_1^i \\
& + \theta^+ \sum_{j=\frac{1}{2}}^{N-\frac{1}{2}} z^{-N+j-\frac{1}{2}} S_j^- \\
& + \theta^+ \sum_{i=0}^{N-1} \sum_{j=\frac{1}{2}}^{N+\frac{1}{2}} z^{-N+j-i-\frac{3}{2}} (p_j^{-,1-,i} L_1 + p_j^{-,2-,i} G_{\frac{1}{2}}^+ G_{\frac{1}{2}}^- + p_{j-\frac{1}{2}}^{-,3--,i} G_{\frac{1}{2}}^+ + p_{j-\frac{1}{2}}^{-,4,i} G_{\frac{1}{2}}^-) L_1^i \\
& + \theta^- \sum_{j=\frac{1}{2}}^{N-\frac{1}{2}} z^{-N+j-\frac{1}{2}} S_j^+ \\
& + \theta^- \sum_{i=0}^{N-1} \sum_{j=\frac{1}{2}}^{N+\frac{1}{2}} z^{-N+j-i-\frac{3}{2}} (p_j^{+,1+,i} L_1 + p_j^{+,2+,i} G_{\frac{1}{2}}^+ G_{\frac{1}{2}}^- + p_{j-\frac{1}{2}}^{+,3,i} G_{\frac{1}{2}}^+ + p_{j-\frac{1}{2}}^{+,4++,i} G_{\frac{1}{2}}^-) L_1^i \\
& + \theta^+ \theta^- \sum_{k=0}^N z^{-N+k-1} \bar{S}_k \\
& + \theta^+ \theta^- \sum_{i=0}^{N-1} \sum_{k=0}^N z^{-N+k-i-2} (\bar{p}_k^{1,i} L_1 + \bar{p}_k^{2,i} G_{\frac{1}{2}}^+ G_{\frac{1}{2}}^- + \bar{p}_{k-\frac{1}{2}}^{3-,i} G_{\frac{1}{2}}^+ + \bar{p}_{k-\frac{1}{2}}^{4+,i} G_{\frac{1}{2}}^-) L_1^i,
\end{aligned} \tag{3.6}$$

where the coefficients^{*h*} $S_k, S_j^-, S_j^+, \bar{S}_k$ and $p_k, p_j^-, p_j^+, \bar{p}_k$ only contain algebra elements in $\mathcal{A}_- \oplus \mathcal{H}$. This allows us to write the coefficients j_n, j_r^+, j_r^- and $\bar{j}_n$ of $\mathcal{J}(Z)$ in terms of f_n, f_r^+, f_r^- and $\bar{f}_n$ using equations (3.4,3.5,3.6).

$$\begin{aligned}
j_n = & \sum_{m=0}^n S_m f_{n-m} + \sum_{i=0}^{N-1} \sum_{m=0}^n (p_m^{1,i} L_1 + p_m^{2,i} G_{\frac{1}{2}}^+ G_{\frac{1}{2}}^- + p_{m-\frac{1}{2}}^{3-,i} G_{\frac{1}{2}}^+ + p_{m-\frac{1}{2}}^{4+,i} G_{\frac{1}{2}}^-) L_1^i f_{n-m+i+1} \\
\bar{j}_n = & \sum_{m=0}^n S_m \bar{f}_{n-m} + \sum_{i=0}^{N-1} \sum_{m=0}^n (p_m^{1,i} L_1 + p_m^{2,i} G_{\frac{1}{2}}^+ G_{\frac{1}{2}}^- + p_{m-\frac{1}{2}}^{3-,i} G_{\frac{1}{2}}^+ + p_{m-\frac{1}{2}}^{4+,i} G_{\frac{1}{2}}^-) L_1^i \bar{f}_{n-m+i+1} \\
& + \sum_{m=0}^n \bar{S}_m f_{n-m} + \sum_{i=0}^{N-1} \sum_{m=0}^n (\bar{p}_m^{1,i} L_1 + \bar{p}_m^{2,i} G_{\frac{1}{2}}^+ G_{\frac{1}{2}}^- + \bar{p}_{m-\frac{1}{2}}^{3-,i} G_{\frac{1}{2}}^+ + \bar{p}_{m-\frac{1}{2}}^{4+,i} G_{\frac{1}{2}}^-) L_1^i f_{n-m+i+1}
\end{aligned}$$

^{*h*}The index coincides with the level of the coefficient.

$$\begin{aligned}
& - \sum_{j=\frac{1}{2}}^{n-\frac{1}{2}} S_j^- f_{n-j}^+ - \sum_{i=0}^{N-1} \sum_{j=\frac{1}{2}}^{n+\frac{1}{2}} (p_j^{-,1-,i} L_1 + p_j^{-,2-,i} G_{\frac{1}{2}}^+ G_{\frac{1}{2}}^- + p_{j-\frac{1}{2}}^{-,3--,i} G_{\frac{1}{2}}^+ + p_{j-\frac{1}{2}}^{-,4,i} G_{\frac{1}{2}}^-) L_1^i f_{n-j+i+1}^+ \\
& + \sum_{j=\frac{1}{2}}^{n-\frac{1}{2}} S_j^+ f_{n-j}^- + \sum_{i=0}^{N-1} \sum_{j=\frac{1}{2}}^{n+\frac{1}{2}} (p_j^{+,1+,i} L_1 + p_j^{+,2+,i} G_{\frac{1}{2}}^+ G_{\frac{1}{2}}^- + p_{j-\frac{1}{2}}^{+,3,i} G_{\frac{1}{2}}^+ + p_{j-\frac{1}{2}}^{+,4++,i} G_{\frac{1}{2}}^-) L_1^i f_{n-j+i+1}^- \\
j_r^- = & \sum_{m=0}^{r-\frac{1}{2}} S_m f_{r-m}^- + \sum_{i=0}^{N-1} \sum_{m=0}^{r+\frac{1}{2}} (p_m^{1,i} L_1 + p_m^{2,i} G_{\frac{1}{2}}^+ G_{\frac{1}{2}}^- + p_{m-\frac{1}{2}}^{3-,i} G_{\frac{1}{2}}^+ + p_{m-\frac{1}{2}}^{4+,i} G_{\frac{1}{2}}^-) L_1^i f_{r-m+i+1}^- \\
& + \sum_{j=\frac{1}{2}}^r S_j^- f_{r-j} + \sum_{i=0}^{N-1} \sum_{j=\frac{1}{2}}^r (p_j^{-,1-,i} L_1 + p_j^{-,2-,i} G_{\frac{1}{2}}^+ G_{\frac{1}{2}}^- + p_{j-\frac{1}{2}}^{-,3--,i} G_{\frac{1}{2}}^+ + p_{j-\frac{1}{2}}^{-,4,i} G_{\frac{1}{2}}^-) L_1^i f_{r-j+i+1}^- \\
j_r^+ = & \sum_{m=0}^{r-\frac{1}{2}} S_m f_{r-m}^+ + \sum_{i=0}^{N-1} \sum_{m=0}^{r+\frac{1}{2}} (p_m^{1,i} L_1 + p_m^{2,i} G_{\frac{1}{2}}^+ G_{\frac{1}{2}}^- + p_{m-\frac{1}{2}}^{3-,i} G_{\frac{1}{2}}^+ + p_{m-\frac{1}{2}}^{4+,i} G_{\frac{1}{2}}^-) L_1^i f_{r-m+i+1}^+ \\
& + \sum_{j=\frac{1}{2}}^r S_j^+ f_{r-j} + \sum_{i=0}^{N-1} \sum_{j=\frac{1}{2}}^r (p_j^{+,1+,i} L_1 + p_j^{+,2+,i} G_{\frac{1}{2}}^+ G_{\frac{1}{2}}^- + p_{j-\frac{1}{2}}^{+,3,i} G_{\frac{1}{2}}^+ + p_{j-\frac{1}{2}}^{+,4++,i} G_{\frac{1}{2}}^-) L_1^i f_{r-j+i+1}^+
\end{aligned} \tag{3.7}$$

We can use the descent equations (2.16) to evaluate the action of L_1 , $G_{\frac{1}{2}}^+$ and $G_{\frac{1}{2}}^-$ on f_n , f_r^+ , f_r^- and $\bar{f}_n$:

$$\begin{aligned}
L_1^{i+1} f_m &= \prod_{j=0}^i (h_1 + L_0 - h_2 + j) f_{m-i-1} \\
L_1^{i+1} f_r^\pm &= \prod_{j=0}^i (h_1 + L_0 - h_2 + j + \frac{1}{2}) f_{r-i-1}^\pm \\
L_1^{i+1} \bar{f}_m &= \prod_{j=0}^i (h_1 + L_0 - h_2 + j + 1) \bar{f}_{m-i-1} \\
&\quad + q_1 \sum_{l=0}^i \prod_{j_1=l+1}^i (h_1 + L_0 - h_2 + j_1 + 1) \prod_{j_2=0}^{l-1} (h_1 + L_0 - h_2 + j_2) f_{m-i-1} \\
G_{\frac{1}{2}}^+ L_1^i f_m &= - \prod_{j=0}^{i-1} (h_1 + L_0 - h_2 + j + \frac{1}{2}) f_{m-i-\frac{1}{2}}^+ \\
G_{\frac{1}{2}}^+ L_1^i f_r^+ &= 0 \\
G_{\frac{1}{2}}^+ L_1^i f_r^- &= - \prod_{j=0}^{i-1} (h_1 + L_0 - h_2 + j + 1) \left((h_1 + L_0 - h_2 + q_1) f_{r-i-\frac{1}{2}}^- + \bar{f}_{r-i-\frac{1}{2}} \right)
\end{aligned}$$

ⁱWe define the empty product $\prod_{i=m}^n \dots$ for $m > n$ to be 1, and the empty sum $\sum_{i=m}^n \dots$ for $m > n$ to vanish.

$$\begin{aligned}
G_{\frac{1}{2}}^+ L_1^i \bar{f}_m &= \left(\prod_{j=0}^{i-1} (h_1 + L_0 - h_2 + j + \frac{3}{2}) (h_1 + L_0 - h_2 + \frac{1}{2} + q_1) \right. \\
&\quad \left. - q_1 \sum_{l=0}^{i-1} \prod_{j_1=l+1}^{i-1} (h_1 + L_0 - h_2 + j_1 + \frac{3}{2}) \prod_{j_2=0}^{l-1} (h_1 + L_0 - h_2 + j_2 + \frac{1}{2}) \right) f_{m-i-\frac{1}{2}}^+ \\
G_{\frac{1}{2}}^- L_1^i f_m &= - \prod_{j=0}^{i-1} (h_1 + L_0 - h_2 + j + \frac{1}{2}) f_{m-i-\frac{1}{2}}^- \\
G_{\frac{1}{2}}^- L_1^i f_r^+ &= - \prod_{j=0}^{i-1} (h_1 + L_0 - h_2 + j + 1) \left((h_1 + L_0 - h_2 - q_1) f_{r-i-\frac{1}{2}} - \bar{f}_{r-i-\frac{1}{2}} \right) \\
G_{\frac{1}{2}}^- L_1^i f_r^- &= 0 \\
G_{\frac{1}{2}}^- L_1^i \bar{f}_m &= - \left(\prod_{j=0}^{i-1} (h_1 + L_0 - h_2 + j + \frac{3}{2}) (h_1 + L_0 - h_2 + \frac{1}{2} - q_1) \right. \\
&\quad \left. + q_1 \sum_{l=0}^{i-1} \prod_{j_1=l+1}^{i-1} (h_1 + L_0 - h_2 + j_1 + \frac{3}{2}) \prod_{j_2=0}^{l-1} (h_1 + L_0 - h_2 + j_2 + \frac{1}{2}) \right) f_{m-i-\frac{1}{2}}^- \\
G_{\frac{1}{2}}^+ G_{\frac{1}{2}}^- L_1^i f_m &= \prod_{j=0}^{i-1} (h_1 + L_0 - h_2 + j + 1) \left((h_1 + L_0 - h_2 + q_1) f_{m-i-1} + \bar{f}_{m-i-1} \right) \\
G_{\frac{1}{2}}^+ G_{\frac{1}{2}}^- L_1^i f_r^+ &= 2 \prod_{j=0}^{i-1} (h_1 + L_0 - h_2 + j + \frac{3}{2}) (h_1 + L_0 - h_2 + \frac{1}{2}) f_{r-i-1}^+ \\
G_{\frac{1}{2}}^+ G_{\frac{1}{2}}^- L_1^i f_r^- &= 0 \\
G_{\frac{1}{2}}^+ G_{\frac{1}{2}}^- L_1^i \bar{f}_m &= \left(\prod_{j=0}^{i-1} (h_1 + L_0 - h_2 + j + 2) (h_1 + L_0 - h_2 + 1 - q_1) \right. \\
&\quad \left. + q_1 \sum_{l=0}^{i-1} \prod_{j_1=l+1}^{i-1} (h_1 + L_0 - h_2 + j_1 + 2) \prod_{j_2=0}^{l-1} (h_1 + L_0 - h_2 + j_2 + 1) \right) \\
&\quad \left((h_1 + L_0 - h_2 + q_1) f_{m-i-1} + \bar{f}_{m-i-1} \right). \tag{3.8}
\end{aligned}$$

As mentioned above we set j_n, j_r^+, j_r^- and $\bar{j}_n$ equal to 0, and solve the equations (3.7) for f_n, f_r^+, f_r^- and $\bar{f}_n$, leading to a set of recursion formulae for them. The recursion procedure splits up in “even” (3.9) and “odd” (3.10) steps which we have to perform alternately.

$$\begin{pmatrix} f_n \\ \bar{f}_n \end{pmatrix} = -\mathcal{S}_{h_j+n, q_j}^{-1} \begin{pmatrix} Y_n^1 \\ Y_n^2 \end{pmatrix} \tag{3.9}$$

$$f_r^+ = -\frac{1}{s_{h_j+r, q_j+1}^+} Y_r^+ \quad f_r^- = -\frac{1}{s_{h_j+r, q_j-1}^-} Y_r^- \tag{3.10}$$

The matrix $\mathcal{S}$ and the coefficients s^+, s^- depend on L_0 and hence on the level n of $f_n, f_n^\pm$ and $\bar{f}_n$. If necessary we shall assign a subscript denoting the level. $\mathcal{S}$ has to be inverted for each

even step; if this fails at a certain level m , we can not define the corresponding j_m and $\bar{j}_m$ to be 0. Similarly we have to divide by s^+ and s^- for each odd step which leads to non trivial $j_r^\pm$ if the division fails:

$$\begin{aligned}
s_{11} &= S_0 + \sum_{i=0}^{N-1} \left\{ p_0^{1,i} \prod_{j=0}^i (h_1 + L_0 - h_2 + j) + p_0^{2,i} \prod_{j=0}^{i-1} (h_1 + L_0 - h_2 + j + 1) \right. \\
&\quad \left. (h_1 + L_0 - h_2 + q_1) \right\} \\
s_{12} &= \sum_{i=0}^{N-1} p_0^{2,i} \prod_{j=0}^{i-1} (h_1 + L_0 - h_2 + j + 1) \\
s_{21} &= \bar{S}_0 + \sum_{i=0}^{N-1} \left\{ q_1 p_0^{1,i} \sum_{l=0}^i \prod_{j_1=l+1}^i (h_1 + L_0 - h_2 + j_1 + 1) \prod_{j_2=0}^{l-1} (h_1 + L_0 - h_2 + j_2) \right. \\
&\quad + p_0^{2,i} \left(\prod_{j=0}^{i-1} (h_1 + L_0 - h_2 + j + 2) (h_1 + L_0 - h_2 + 1 - q_1) + q_1 \right. \\
&\quad \left. \sum_{l=0}^{i-1} \prod_{j_1=l+1}^{i-1} (h_1 + L_0 - h_2 + j_1 + 2) \prod_{j_2=0}^{l-1} (h_1 + L_0 - h_2 + j_2 + 1) \right) (h_1 + L_0 - h_2 + q_1) \\
&\quad + \bar{p}_0^{1,i} \prod_{j=0}^i (h_1 + L_0 - h_2 + j) + \bar{p}_0^{2,i} \prod_{j=0}^{i-1} (h_1 + L_0 - h_2 + j + 1) (h_1 + L_0 - h_2 + q_1) \\
&\quad \left. + \prod_{j=0}^{i-1} (h_1 + L_0 - h_2 + j + 1) (p_0^{-,4,i} (h_1 + L_0 - h_2 - q_1) - p_0^{+,3,i} (h_1 + L_0 - h_2 + q_1)) \right\} \\
s_{22} &= S_0 + \sum_{i=0}^{N-1} \left\{ p_0^{1,i} \prod_{j=0}^i (h_1 + L_0 - h_2 + j + 1) + p_0^{2,i} \left(\prod_{j=0}^{i-1} (h_1 + L_0 - h_2 + j + 2) \right. \right. \\
&\quad \left. (h_1 + L_0 - h_2 + 1 - q_1) + q_1 \sum_{l=0}^{i-1} \prod_{j_1=l+1}^{i-1} (h_1 + L_0 - h_2 + j_1 + 2) \right. \\
&\quad \left. \prod_{j_2=0}^{l-1} (h_1 + L_0 - h_2 + j_2 + 1) \right) + \bar{p}_0^{2,i} \prod_{j=0}^{i-1} (h_1 + L_0 - h_2 + j + 1) \\
&\quad \left. - \prod_{j=0}^{i-1} (h_1 + L_0 - h_2 + j + 1) (p_0^{-,4,i} + p_0^{+,3,i}) \right\} \tag{3.11}
\end{aligned}$$

$$\mathcal{S} = \begin{pmatrix} s_{11} & s_{12} \\ s_{21} & s_{22} \end{pmatrix} \tag{3.12}$$

$$\begin{aligned}
s^+ &= S_0 + \sum_{i=0}^{N-1} \left\{ p_0^{1,i} \prod_{j=0}^i (h_1 + L_0 - h_2 + j + \frac{1}{2}) + 2p_0^{2,i} \prod_{j=0}^{i-1} (h_1 + L_0 - h_2 + j + \frac{3}{2}) \right. \\
&\quad \left. (h_1 + L_0 - h_2 + \frac{1}{2}) - p_0^{+,3,i} \prod_{j=0}^{i-1} (h_1 + L_0 - h_2 + j + \frac{1}{2}) \right\}
\end{aligned}$$

$$s^- = S_0 + \sum_{i=0}^{N-1} \left\{ p_0^{1,i} \prod_{j=0}^i (h_1 + L_0 - h_2 + j + \frac{1}{2}) - p_0^{-,4,i} \prod_{j=0}^{i-1} (h_1 + L_0 - h_2 + j + \frac{1}{2}) \right\}. \quad (3.13)$$

The expressions for Y_n^1 and Y_n^2 depend on $f_m, f_s^\pm$ and $\bar{f}_m$ ($m, s < n$), similarly for $Y_r^\pm$, so that (3.9, 3.10) really defines a recursion. For convenience we give the following expressions for Y_n^1, Y_n^2 and $Y_r^\pm$ in which we still have to insert equations (3.8):

$$\begin{aligned} Y_n^1 &= \sum_{m=1}^n S_m f_{n-m} + \sum_{i=0}^{N-1} \sum_{m=1}^n (p_m^{1,i} L_1 + p_m^{2,i} G_{\frac{1}{2}}^+ G_{\frac{1}{2}}^- + p_{m-\frac{1}{2}}^{3-,i} G_{\frac{1}{2}}^+ + p_{m-\frac{1}{2}}^{4+,i} G_{\frac{1}{2}}^-) L_1^i f_{n-m+i+1} \\ Y_n^2 &= \sum_{m=1}^n S_m \bar{f}_{n-m} + \sum_{i=0}^{N-1} \sum_{m=1}^n (p_m^{1,i} L_1 + p_m^{2,i} G_{\frac{1}{2}}^+ G_{\frac{1}{2}}^- + p_{m-\frac{1}{2}}^{3-,i} G_{\frac{1}{2}}^+ + p_{m-\frac{1}{2}}^{4+,i} G_{\frac{1}{2}}^-) L_1^i \bar{f}_{n-m+i+1} \\ &\quad + \sum_{m=1}^n \bar{S}_m f_{n-m} + \sum_{i=0}^{N-1} \sum_{m=1}^n (\bar{p}_m^{1,i} L_1 + \bar{p}_m^{2,i} G_{\frac{1}{2}}^+ G_{\frac{1}{2}}^- + \bar{p}_{m-\frac{1}{2}}^{3-,i} G_{\frac{1}{2}}^+ + \bar{p}_{m-\frac{1}{2}}^{4+,i} G_{\frac{1}{2}}^-) L_1^i f_{n-m+i+1} \\ &\quad - \sum_{j=\frac{1}{2}}^{n-\frac{1}{2}} S_j^+ f_{n-j}^+ - \sum_{i=0}^{N-1} \sum_{j=\frac{1}{2}}^{n-\frac{1}{2}} (p_j^{-,1-,i} L_1 + p_j^{-,2-,i} G_{\frac{1}{2}}^+ G_{\frac{1}{2}}^-) L_1^i f_{n-j+i+1}^+ \\ &\quad - \sum_{i=0}^{N-1} \sum_{j=\frac{3}{2}}^{n+\frac{1}{2}} (p_{j-\frac{1}{2}}^{-,3--,i} G_{\frac{1}{2}}^+ + p_{j-\frac{1}{2}}^{-,4,i} G_{\frac{1}{2}}^-) L_1^i f_{n-j+i+1}^+ \\ &\quad + \sum_{j=\frac{1}{2}}^{n-\frac{1}{2}} S_j^+ f_{n-j}^- + \sum_{i=0}^{N-1} \sum_{j=\frac{1}{2}}^{n-\frac{1}{2}} (p_j^{+,1+,i} L_1 + p_j^{+,2+,i} G_{\frac{1}{2}}^+ G_{\frac{1}{2}}^-) L_1^i f_{n-j+i+1}^- \\ &\quad + \sum_{i=0}^{N-1} \sum_{j=\frac{3}{2}}^{n+\frac{1}{2}} (p_{j-\frac{1}{2}}^{+,3,i} G_{\frac{1}{2}}^+ + p_{j-\frac{1}{2}}^{+,4+,i} G_{\frac{1}{2}}^-) L_1^i f_{n-j+i+1}^- \end{aligned} \quad (3.14)$$

$$\begin{aligned} Y_r^- &= \sum_{m=1}^{r-\frac{1}{2}} S_m f_{r-m}^- + \sum_{i=0}^{N-1} \sum_{m=1}^{r+\frac{1}{2}} (p_m^{1,i} L_1 + p_m^{2,i} G_{\frac{1}{2}}^+ G_{\frac{1}{2}}^- + p_{m-\frac{1}{2}}^{3-,i} G_{\frac{1}{2}}^+ + p_{m-\frac{1}{2}}^{4+,i} G_{\frac{1}{2}}^-) L_1^i f_{r-m+i+1}^- \\ &\quad + \sum_{j=\frac{1}{2}}^r S_j^- f_{r-j}^- + \sum_{i=0}^{N-1} \sum_{j=\frac{1}{2}}^r (p_j^{-,1-,i} L_1 + p_j^{-,2-,i} G_{\frac{1}{2}}^+ G_{\frac{1}{2}}^-) L_1^i f_{r-j+i+1}^- \\ &\quad + \sum_{i=0}^{N-1} \sum_{j=\frac{3}{2}}^r (p_{j-\frac{1}{2}}^{-,3--,i} G_{\frac{1}{2}}^+ + p_{j-\frac{1}{2}}^{-,4,i} G_{\frac{1}{2}}^-) L_1^i f_{r-j+i+1}^- \\ Y_r^+ &= \sum_{m=1}^{r-\frac{1}{2}} S_m f_{r-m}^+ + \sum_{i=0}^{N-1} \sum_{m=1}^{r+\frac{1}{2}} (p_m^{1,i} L_1 + p_m^{2,i} G_{\frac{1}{2}}^+ G_{\frac{1}{2}}^- + p_{m-\frac{1}{2}}^{3-,i} G_{\frac{1}{2}}^+ + p_{m-\frac{1}{2}}^{4+,i} G_{\frac{1}{2}}^-) L_1^i f_{r-m+i+1}^+ \\ &\quad + \sum_{j=\frac{1}{2}}^r S_j^+ f_{r-j}^+ + \sum_{i=0}^{N-1} \sum_{j=\frac{1}{2}}^r (p_j^{+,1+,i} L_1 + p_j^{+,2+,i} G_{\frac{1}{2}}^+ G_{\frac{1}{2}}^-) L_1^i f_{r-j+i+1}^+ \end{aligned}$$

$$+ \sum_{i=0}^{N-1} \sum_{j=\frac{3}{2}}^r (p_{j-\frac{1}{2}}^{+,3,i} G_{\frac{1}{2}}^+ + p_{j-\frac{1}{2}}^{+,4^{++},i} G_{\frac{1}{2}}^-) L_1^i f_{r-j+i+1} \quad (3.15)$$

The initial conditions for the recursion can be found easily by considering that we have no singular vectors at level 0. Hence, in order to decouple the singular vector $\mathcal{N}$ we necessarily need j_0 and $\bar{j}_0$ to vanish. The only terms contributing to them are f_0 and $\bar{f}_0$, which are both multiples of the highest weight vector: $f_0 = c_f |h_j, q_j\rangle$, $\bar{f}_0 = c_{\bar{f}} |h_j, q_j\rangle$. Using equations (3.7) we find:

$$\begin{pmatrix} 0 \\ 0 \end{pmatrix} = \mathcal{S}_{h_j, q_j} \begin{pmatrix} c_f \\ c_{\bar{f}} \end{pmatrix}, \quad (3.16)$$

where the matrix $\mathcal{S}_{h_j, q_j}$ is taken at level 0: $L_0 = h_j$ and $T_0 = q_j$. According to (3.16), $(c_f, c_{\bar{f}})^T$ has to be an eigenvector of $\mathcal{S}_{h_j, q_j}$ with respect to the eigenvalue 0. This determines restrictions on the allowed fusions to be the ones where $\mathcal{S}_{h_j, q_j}$ has an eigenvalue 0.

If the recursion breaks down at a certain level r because s_{h_j+r, q_j+1}^+ or s_{h_j+r, q_j-1}^- happens to be 0 then we can use (3.7) to compute j_r^+ or j_r^- respectively which will be singular. If at a certain level m , $\mathcal{S}_{h_j+m, q_j}$ happens to be singular, we get a singular vector by using (3.7) to compute the combination of j_m and $\bar{j}_m$ which corresponds to the eigenvalue 0 and hence does not involve the unknown terms f_m and $\bar{f}_m$. This completes the recursion procedure. In the following section we will outline an induction proof for it.

Although the recursion formulae look rather complicated, in the actual calculation they turn out to simplify rather drastically.

3.2. Proof of the recursion procedure

We shall now forget about the heuristic arguments used in the section above and summarise the recursion procedure: we use as initial conditions f_0 and $\bar{f}_0$ according to equation (3.16); we perform odd and even recursion steps for $f_n, \bar{f}_n, f_r^\pm$ alternately according to equations (3.9, 3.10) until the procedure breaks down at level m because $\mathcal{S}_{h_j+m, q_j}, s_{h_j+m, q_j+1}^+$ or s_{h_j+m, q_j-1}^- is not invertible any more. The coefficients $f_n, \bar{f}_n, f_r^\pm$ ($n, r < m$) obtained in this way satisfy the descent equations (2.16), the $j_n, \bar{j}_n, j_r^\pm$ ($n, r < m$) vanish and respectively $j_m^\pm$ or the $j_m, \bar{j}_m$ combination corresponding to the $\mathcal{S}_{h_j+m, q_j}$ eigenvalue 0 at level m is a singular vector or vanishes completely.

We shall prove this by induction on the level n . For $n = 0$ we have chosen $j_0 = \bar{j}_0$ equals 0 by construction of f_0 and $\bar{f}_0$ which are multiples of the highest weight vector $|h_j, q_j\rangle$ and hence satisfy the descent equations (2.16). As induction hypothesis we shall assume now:

$$\begin{aligned} \det(\mathcal{S}_{h_j, q_j}) &= 0 \\ \det(\mathcal{S}_{h_j+k, q_j}) &\neq 0 \quad , \quad 0 < k < n \\ s_{h_j+r, q_j \pm 1}^\pm &\neq 0 \quad , \quad 0 < r < n \end{aligned} \quad (3.17)$$

and the descent equations are satisfied by $f_k, f_r^\pm$ and $\bar{f}_k$ for $0 \leq k < n$, $0 < r < n$.

Assume n is integral^j. We show that either $\det(\mathcal{S}_{h_j+n, q_j}) \neq 0$ which will imply the descent equations for f_n and $\bar{f}_n$, or $\det(\mathcal{S}_{h_j+n, q_j}) = 0$ which will imply the construction of a singular vector at level n .

As a first step we want to translate $G_{\frac{1}{2}}^\pm$ and L_1 back to superderivative language. If we use the notation

$$\begin{aligned}\gamma_1 &= 2h_1 z + z\theta^+ \frac{\partial}{\partial\theta^+} + z\theta^- \frac{\partial}{\partial\theta^-} + z^2 \partial_z + q_1 \theta^+ \theta^- \\ \gamma_\pm &= 2h_1 \theta^\pm - z \frac{\partial}{\partial\theta^\mp} + \theta^\pm z \partial_z \pm \theta^+ \theta^- \frac{\partial}{\partial\theta^\mp} \pm q_1 \theta^\pm,\end{aligned}\quad (3.18)$$

we find:

$$\begin{aligned}L_1^{i+1} \Phi_{h_1, q_1}(Z) |h_2, q_2\rangle &= \gamma_1^{i+1} \Phi_{h_1, q_1}(Z) |h_2, q_2\rangle \\ G_{\frac{1}{2}}^\pm L_1^i \Phi_{h_1, q_1}(Z) |h_2, q_2\rangle &= \gamma_1^i \gamma_\pm \Phi_{h_1, q_1}(Z) |h_2, q_2\rangle \\ G_{\frac{1}{2}}^+ G_{\frac{1}{2}}^- L_1^i \Phi_{h_1, q_1}(Z) |h_2, q_2\rangle &= \gamma_1^i \gamma_- \gamma_+ \Phi_{h_1, q_1}(Z) |h_2, q_2\rangle.\end{aligned}\quad (3.19)$$

The left ideal which annihilates the highest weight state $|h_2, q_2\rangle$ is equally well spanned by $\{(l_m + h_2 z \delta_{m,1}), (t_m + q_2 z \delta_{m,1}), g_r^\pm, G_{\frac{1}{2}}^\pm, (L_0 - h_2), (T_0 - q_2); m \in \mathbb{N}, r \in \mathbb{N} + \frac{1}{2}\}$, using (2.12). Since $\mathcal{N} |h_2, q_2\rangle$ is singular, the above left ideal annihilates this vector. Hence, and by considering the commutation relations we get:

$$\begin{aligned}G_{\frac{1}{2}}^+ \mathcal{N} &= \mathcal{N} G_{\frac{1}{2}}^+ + \mathcal{N}^{1,+} (L_0 - h_2) + \mathcal{N}^{2,+} (T_0 - q_2) \\ G_{\frac{1}{2}}^- \mathcal{N} &= \mathcal{N} G_{\frac{1}{2}}^- + \mathcal{N}^{1,-} (L_0 - h_2) + \mathcal{N}^{2,-} (T_0 - q_2) \\ (t_1 + q_2 z) \mathcal{N} &= \mathcal{N} (t_1 + q_2 z) + \mathcal{N}'^- G_{\frac{1}{2}}^+ + \mathcal{N}'^+ G_{\frac{1}{2}}^- + \mathcal{N}'^1 (L_0 - h_2) + \mathcal{N}'^2 (T_0 - q_2).\end{aligned}\quad (3.20)$$

It is easy to see that $\sigma(L_0 - h_2) \mathcal{F}(Z) = \sigma(T_0 - q_2) \mathcal{F}(Z) = 0$. We apply the representation σ (2.10) to (3.20). For simplicity we shall only be interested in the contributions involving no $\theta^\pm$ terms, however, the following considerations can equally well be done for them. We take the expressions (2.10, 3.6) for $\sigma(G_{\frac{1}{2}}^+) \sigma(\mathcal{N}) \mathcal{F}(Z)$ and use equations (3.18) to replace algebra generators by superderivatives. By considering the commutation relations we interchange $\sigma(G_{\frac{1}{2}}^+)$ and $\sigma(\mathcal{N})$ and obtain a form according to (3.20):^k

$$\begin{aligned}\sigma(G_{\frac{1}{2}}^+) \sigma(\mathcal{N}) \mathcal{F}(Z) &= \left\{ (G_{\frac{1}{2}}^+ + z \frac{\partial}{\partial\theta^-}) \left(\frac{1}{z^N} S_0 + \sum_{i=0}^{N-1} \frac{1}{z^{N+i+1}} (p_0^{1,i} L_1 + p_0^{2,i} G_{\frac{1}{2}}^+ G_{\frac{1}{2}}^-) L_1^i + \dots \right) \right. \\ &\quad \left. + \mathcal{O}(\theta^\pm) + \mathcal{O}(\theta^+ \theta^-) \right\} \mathcal{F}(Z) \\ &= \left\{ \left(\frac{1}{z^N} S_{0; L_0 + \frac{1}{2}, T_0 - 1} + \sum_{i=0}^{N-1} \frac{1}{z^{N+i+1}} (p_{0; L_0 + \frac{1}{2}, T_0 - 1}^{1,i} \gamma_1^{i+1} \right. \right.\end{aligned}$$

^jSimilar arguments can be used if n is half integral.

^k $S_{0; L_0 + \frac{1}{2}, T_0 - 1}$ denotes S_0 in which we have replaced L_0 by $L_0 + \frac{1}{2}$ and T_0 by $T_0 - 1$ etc.

$$\begin{aligned}
& + p_{0;L_0+\frac{1}{2},T_0-1}^{2,i} \gamma_1^i \gamma_- \gamma_+ + \dots)(G_{\frac{1}{2}}^+ + z \frac{\partial}{\partial \theta^-}) \\
& + \mathcal{O}(\theta^\pm) + \mathcal{O}(\theta^+ \theta^-) \} \mathcal{F}(Z).
\end{aligned} \tag{3.21}$$

Hence, using the induction hypothesis $G_{\frac{1}{2}}^+ f_k + f_{k-\frac{1}{2}}^+ = 0$ for $k < n$:

$$\begin{aligned}
\sigma(G_{\frac{1}{2}}^+) \mathcal{J}(Z) &= \left\{ \frac{1}{z^N} S_{0;L_0+\frac{1}{2},T_0-1} + \sum_{i=0}^{N-1} \frac{1}{z^{N+i+1}} (p_{0;L_0+\frac{1}{2},T_0-1}^{1,i} \gamma_1^{i+1} \right. \\
& \quad \left. + p_{0;L_0+\frac{1}{2},T_0-1}^{2,i} \gamma_1^i \gamma_- \gamma_+ + \dots) \right\} \left(\sum_{k \geq n} z^{h_j-h_1-h_2+k} (G_{\frac{1}{2}}^+ f_k + f_{k-\frac{1}{2}}^+) \right) \\
& \quad + \mathcal{O}(\theta^\pm) + \mathcal{O}(\theta^+ \theta^-).
\end{aligned} \tag{3.22}$$

The action of γ_1^{i+1} and $\gamma_\pm$ on $\sum_{k \geq n} z^{h_j-h_1-h_2+k} (G_{\frac{1}{2}}^+ f_k + f_{k-\frac{1}{2}}^+)$ reproduces the coefficients of $p_0^{1,i}$ and $p_0^{2,i}$ in equations (3.11). Comparing coefficients and doing the same for the $\theta^\pm$ and $\theta^+ \theta^-$ components, we find:

$$\begin{pmatrix} G_{\frac{1}{2}}^+ j_n \\ G_{\frac{1}{2}}^+ \bar{j}_n \end{pmatrix} = \mathcal{S}_{h_j+n,q_j} \begin{pmatrix} G_{\frac{1}{2}}^+ f_n + f_{n-\frac{1}{2}}^+ \\ G_{\frac{1}{2}}^+ \bar{f}_n - (h_1 + L_0 - h_2 + \frac{1}{2} + q_1) f_{n-\frac{1}{2}}^+ \end{pmatrix} \tag{3.23}$$

Suppose $\det \mathcal{S}$ at level n is non trivial, we have j_n and $\bar{j}_n$ equal to 0 due to the definition of f_n and $\bar{f}_n$. Hence, $G_{\frac{1}{2}}^+ f_n = -f_{n-\frac{1}{2}}^+$ and $G_{\frac{1}{2}}^+ \bar{f}_n = (h_1 + L_0 - h_2 + \frac{1}{2} + q_1) f_{n-\frac{1}{2}}^+$. However, if $\det \mathcal{S}_{h_j+n,q_j}$ at level n does vanish, we multiply (3.23) by the inverse basis transformation $\mathcal{T}$, which transforms $\mathcal{S}_{h_j+n,q_j}$ in its Jordan normal form:

$$\mathcal{T}^{-1} \begin{pmatrix} G_{\frac{1}{2}}^+ j_n \\ G_{\frac{1}{2}}^+ \bar{j}_n \end{pmatrix} = \begin{pmatrix} 0 & 0 \\ * & * \end{pmatrix} \mathcal{T}^{-1} \begin{pmatrix} G_{\frac{1}{2}}^+ f_n + f_{n-\frac{1}{2}}^+ \\ G_{\frac{1}{2}}^+ \bar{f}_n - (h_1 + L_0 - h_2 + \frac{1}{2} + q_1) f_{n-\frac{1}{2}}^+ \end{pmatrix} \tag{3.24}$$

Hence, we find for the first component of $\mathcal{T}^{-1} \begin{pmatrix} j_n \\ \bar{j}_n \end{pmatrix}$, Ψ say, that $G_{\frac{1}{2}}^+ \Psi = 0$.

Similar considerations for $G_{\frac{1}{2}}^-$ and $t_1 + q_2 z$ lead to descent equations for $G_{\frac{1}{2}}^-$ and T_1 , or to $G_{\frac{1}{2}}^+ \Psi = 0$ and $T_1 \Psi = 0$ depending whether $\mathcal{S}_{h_j+n,q_j}$ is invertible or not. Since $G_{\frac{1}{2}}^+$, $G_{\frac{1}{2}}^-$ and T_1 generate $\mathcal{A}_+$ this is already sufficient.

The above considerations turn out to be even easier if n is taken to be positive, half integral, and they lead to the same result depending whether s_{h_j+n,q_j+1}^+ or s_{h_j+n,q_j-1}^- is invertible or not. Whichever fails to be invertible leads to a singular vector j_n^+ or j_n^- respectively, exactly in the same way as for Ψ . This completes the proof^l.

^lThis does not exclude that the recursively computed vector is identical to 0. However, the non triviality can usually be seen by looking at the coefficients of the vector explicitly.

4. Explicit formulae for selected singular vectors

4.1. Positive charged singular vectors

We start off using the singular vector (2.19) of the representation generated by the field $\Phi_{h_{1,2},q'}$ and investigate possible fusions with fields $\Phi_{h_k,q}$ to give fields $\Phi_{h_{\tilde{k}},\tilde{q}}$. Factorising the recursion determinant (3.12) at level 0, we find as allowed fusions:

$$\Phi_{h_{-k},-1-q} \otimes \Phi_{h_{1,2},1+2q} \rightarrow \Phi_{h_k,q} \quad (4.1)$$

$$\Phi_{h_{-k},1-q} \otimes \Phi_{h_{1,2},-1+2q} \rightarrow \Phi_{h_k,q} \quad (4.2)$$

Using either of these two fusions and looking at the recursion formulae shows very easily that if we take the fusion (4.1) the procedure breaks down for positive k , whilst for fusion (4.2) it breaks down for negative k . This allows us to give explicit expressions for all charged singular vectors, which will be done in this and the following section.

We first consider the fusion (4.1), and put k positive, half integral. The recursion formulae given in section 3.1. shows that the only combinations of generators which do appear are

$$\mathcal{L}_{-1}^+ = \frac{1}{2} \frac{t}{q+1} L_{-1} + \frac{1}{4} \frac{t}{q(q+1)} G_{-\frac{1}{2}}^+ G_{-\frac{1}{2}}^- + T_{-1},$$

as well as $G_{-\frac{1}{2}}^+$ and $G_{-\frac{1}{2}}^-$.

Since the recursion formulae involve in this case only terms which differ in level at most by one, we can express the recursion procedure as a product of four by four matrices. We only have to make sure to choose the right initial conditions and finally to project out the singular vector correctly from the four-component vector $(f_{k-\frac{1}{2}}, \bar{f}_{k-\frac{1}{2}}, f_{k-1}^-, f_{k-1}^+)^T$.

The “even” recursion step is given by

$$\mathcal{E}^+(n) = \begin{pmatrix} e_{1,1}^+ \mathcal{L}_{-1}^+ & e_{1,2}^+ \mathcal{L}_{-1}^+ & e_{1,3}^+ G_{-\frac{1}{2}}^+ & e_{1,4}^+ G_{-\frac{1}{2}}^- \\ e_{2,1}^+ \mathcal{L}_{-1}^+ & e_{2,2}^+ \mathcal{L}_{-1}^+ & e_{2,3}^+ G_{-\frac{1}{2}}^+ & e_{2,4}^+ G_{-\frac{1}{2}}^- \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}, \quad \begin{pmatrix} f_n \\ \bar{f}_n \\ f_{n-\frac{1}{2}}^- \\ f_{n-\frac{1}{2}}^+ \end{pmatrix} = \mathcal{E}^+(n) \begin{pmatrix} f_{n-1} \\ \bar{f}_{n-1} \\ f_{n-1}^- \\ f_{n-1}^+ \end{pmatrix}, \quad (4.3)$$

where the matrix entries are

$$\begin{aligned} e_{1,1}^+ &= \frac{t(2q+1)(2(k-n)+1) + 4q(q+1)}{2nt(t(2k-n) + 2(q+1))} \\ e_{1,2}^+ &= -\frac{1}{n(t(2k-n) + 2(q+1))} \\ e_{1,3}^+ &= \frac{t(2(k-n)+1)}{4nq(t(2k-n) + 2(q+1))} \\ e_{1,4}^+ &= -\frac{t(2(k-n)+1) + 4(q+1)}{4n(q+1)(t(2k-n) + 2(q+1))} \end{aligned}$$

$$\begin{aligned}
e_{2,1}^+ &= -\frac{4qt(q+1)(4k^2t+4n-t+4kq)+4t^2(k-n)^2+16q(q+1)^2(kt+q)-t^2}{4nt^2(t(2k-n)+2(q+1))} \\
e_{2,2}^+ &= \frac{t(2q+1)(2(k-n)-1)+4q(q+1)}{2nt(t(2k-n)+2(q+1))} \\
e_{2,3}^+ &= -\frac{t(2q+1)(4k^2-1)+4q(q+1)(2(k+n)+1)-4nt(2k-n)}{8nq(t(2k-n)+2(q+1))} \\
e_{2,4}^+ &= \frac{4nt^2(2k-n)+t^2(2q+1)(4k^2-1)+8t(q+1)(q(3k-n)+k+n)+4(4q-t)(q+1)^2}{8nt(q+1)(t(2k-n)+2(q+1))} \quad .
\end{aligned}$$

The “odd” recursion step can be found to be

$$\mathcal{T}^+(r) = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ \frac{2q(q+1)-t(k-r)}{2(q+1)(t(k-r)+2q)}G_{-\frac{1}{2}}^- & \frac{t}{2(q+1)(t(k-r)+2q)}G_{-\frac{1}{2}}^- & -\frac{2q}{t(k-r)+2q}\mathcal{L}_{-1}^+ & 0 \\ -\frac{2q(q+1)-t(k-r)}{2qt(k-r)}G_{-\frac{1}{2}}^+ & \frac{1}{2q(k-r)}G_{-\frac{1}{2}}^+ & 0 & -\frac{2(q+1)}{t(k-r)}\mathcal{L}_{-1}^+ \end{pmatrix} \quad (4.4)$$

$$\begin{pmatrix} \frac{f_{r-\frac{1}{2}}}{\bar{f}_{r-\frac{1}{2}}} \\ f_r^- \\ f_r^+ \end{pmatrix} = \mathcal{T}^+(r) \begin{pmatrix} \frac{f_{r-\frac{1}{2}}}{\bar{f}_{r-\frac{1}{2}}} \\ f_{r-1}^- \\ f_{r-1}^+ \end{pmatrix}.$$

Finally, we find as initial conditions

$$\begin{pmatrix} \frac{f_0}{\bar{f}_0} \\ f_{-\frac{1}{2}}^- \\ f_{-\frac{1}{2}}^+ \end{pmatrix} = \psi_0^+ = \begin{pmatrix} -2t|h_k, q\rangle \\ (4q(q+1) + (2q+1)(2k-1)t)|h_k, q\rangle \\ 0 \\ 0 \end{pmatrix}, \quad (4.5)$$

where $|h_k, q\rangle$ denotes the highest weight vector, and we take

$$\Psi_k = \underbrace{\left(\frac{1}{2}G_{-\frac{1}{2}}^+, -\frac{1}{4}\frac{t}{q(q+1)}G_{-\frac{1}{2}}^+, 0, \mathcal{L}_{-1}^+ \right)}_{=\mathcal{W}^+} \begin{pmatrix} \frac{f_{k-\frac{1}{2}}}{\bar{f}_{k-\frac{1}{2}}} \\ f_{k-1}^- \\ f_{k-1}^+ \end{pmatrix}. \quad (4.6)$$

Altogether, we can write the $+1$ charged singular vector at level k in the Verma module $\mathcal{V}_{h_k, q}$ with k in $\mathbb{N}_0 + \frac{1}{2}$ in the form

$$\Psi_k = \mathcal{W}^+ \mathcal{E}^+(k - \frac{1}{2}) \mathcal{T}^+(k-1) \mathcal{E}^+(k - \frac{3}{2}) \mathcal{T}^+(k-2) \dots \mathcal{E}^+(1) \mathcal{T}^+(\frac{1}{2}) \psi_0^+. \quad (4.7)$$

4.2. Negative charged singular vectors

Using the fusion (4.2) allows us to compute the singular vectors of the Verma modules $\mathcal{V}_{h_k, q}$ for k negative, half integral. Similarly to the previous section we find

$$\mathcal{L}_{-1}^{-} = \frac{t}{2q} L_{-1} + \frac{1}{4} \frac{t}{q(q-1)} G_{-\frac{1}{2}}^{+} G_{-\frac{1}{2}}^{-} + T_{-1},$$

$$\mathcal{E}^{-}(n) = \begin{pmatrix} e_{1,1}^{-} \mathcal{L}_{-1}^{-} & e_{1,2}^{-} \mathcal{L}_{-1}^{-} & e_{1,3}^{-} G_{-\frac{1}{2}}^{+} & e_{1,4}^{-} G_{-\frac{1}{2}}^{-} \\ e_{2,1}^{-} \mathcal{L}_{-1}^{-} & e_{2,2}^{-} \mathcal{L}_{-1}^{-} & e_{2,3}^{-} G_{-\frac{1}{2}}^{+} & e_{2,4}^{-} G_{-\frac{1}{2}}^{-} \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}, \quad \begin{pmatrix} f_n \\ \bar{f}_n \\ f_{n-\frac{1}{2}}^{-} \\ f_{n-\frac{1}{2}}^{+} \end{pmatrix} = \mathcal{E}^{-}(n) \begin{pmatrix} f_{n-1} \\ \bar{f}_{n-1} \\ f_{n-\frac{1}{2}}^{-} \\ f_{n-\frac{1}{2}}^{+} \end{pmatrix}, \quad (4.8)$$

$$\begin{aligned} e_{1,1}^{-} &= \frac{t(2q-1)(2(k+n)-1) + 4q(q-1)}{2nt(t(2k+n) + 2(q-1))} \\ e_{1,2}^{-} &= \frac{1}{n(t(2k+n) + 2(q-1))} \\ e_{1,3}^{-} &= \frac{t(2(k+n)-1) + 4(q-1)}{4n(q-1)(t(2k+n) + 2(q-1))} \\ e_{1,4}^{-} &= -\frac{t(2(k+n)-1)}{4nq(t(2k+n) + 2(q-1))} \\ e_{2,1}^{-} &= \frac{4qt(q-1)(4k^2t + 4n - t + 4kq) + 4t^2(k+n)^2 + 16q(q-1)^2(kt+q) - t^2}{4nt^2(t(2k+n) + 2(q-1))} \\ e_{2,2}^{-} &= \frac{t(2q-1)(2(k+n)+1) + 4q(q-1)}{2nt(t(2k+n) + 2(q-1))} \\ e_{2,3}^{-} &= \frac{4nt^2(2k+n) + t^2(2q-1)(4k^2-1) + 8t(q-1)(q(3k+n) - k+n) + 4(4q+t)(q-1)^2}{8nt(q-1)(t(2k+n) + 2(q-1))} \\ e_{2,4}^{-} &= -\frac{(t(2q-1)(4k^2-1) + 4q(q-1)(2(k-n)-1) - 4nt(2k+n))}{8nq(t(2k+n) + 2(q-1))}, \end{aligned}$$

$$T^{-}(r) = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ -\frac{2q(q-1)+t(k+r)}{2qt(k+r)} G_{-\frac{1}{2}}^{-} & -\frac{1}{2q(k+r)} G_{-\frac{1}{2}}^{-} & \frac{2(q-1)}{t(k+r)} \mathcal{L}_{-1}^{-} & 0 \\ \frac{2q(q-1)+t(k+r)}{2(q-1)(t(k+r)+2q)} G_{-\frac{1}{2}}^{+} & -\frac{t}{2(q-1)(t(k+r)+2q)} G_{-\frac{1}{2}}^{+} & 0 & \frac{2q}{t(k+r)+2q} \mathcal{L}_{-1}^{-} \end{pmatrix} \quad (4.9)$$

$$\begin{pmatrix} f_0 \\ \bar{f}_0 \\ f_{-\frac{1}{2}}^{-} \\ f_{-\frac{1}{2}}^{+} \end{pmatrix} = \psi_0^{-} = \begin{pmatrix} 2t |h_k, q\rangle \\ (4q(q-1) + (2q-1)(2k+1)t) |h_k, q\rangle \\ 0 \\ 0 \end{pmatrix}, \quad (4.10)$$

$$\Psi_k = \underbrace{\left(-\frac{1}{2}G_{-\frac{1}{2}}^-, -\frac{1}{4}\frac{t}{q(q-1)}G_{-\frac{1}{2}}^-, \mathcal{L}_{-1}^-, 0\right)}_{=\mathcal{W}^-} \begin{pmatrix} f_{k-\frac{1}{2}} \\ \bar{f}_{k-\frac{1}{2}} \\ f_{k-1}^- \\ f_{k-1}^+ \end{pmatrix}. \quad (4.11)$$

And finally, we can write down the -1 charged singular vector in the Verma module $\mathcal{V}_{h_k, q}$ ($-k \in \mathbb{N}_0 + \frac{1}{2}$) at level $|k|$:

$$\Psi_k = \mathcal{W}^- \mathcal{E}^- (|k| - \frac{1}{2}) \mathcal{T}^- (|k| - 1) \mathcal{E}^- (|k| - \frac{3}{2}) \mathcal{T}^- (|k| - 2) \dots \mathcal{E}^- (1) \mathcal{T}^- (\frac{1}{2}) \psi_0^-. \quad (4.12)$$

4.3. Uncharged $(r, 2)$ singular vectors

In this section we use the recursion in order to produce uncharged singular vectors $\Psi_{r, s}$. Factorising the recursion determinant at level 0 gives the allowed fusions

$$\Phi_{h_{r, s, q-\tilde{q}}} \otimes \Phi_{h_{1, 2, \tilde{q}}} \rightarrow \Phi_{h_{r, s-2, q}} + \Phi_{h_{r, s+2, q}}. \quad (4.13)$$

Factorising the recursion matrix at the level where we are looking for the singular vector tells us that the recursion will break down for the cases we consider if and only if $s = 0$. However, this is not at all surprising since we expect to find another freedom in solving the descent equations if the Verma module corresponding to $\Phi_{h_{r, s, q-\tilde{q}}}$ is reducible. In such a case we would have to use information about the singular vector of $\mathcal{V}_{h_{r, s, q-\tilde{q}}}$ as well, as suggested by Bauer et al.¹⁵ for the Virasoro case. Putting $s = 0$ allows us to compute the singular vectors $\Psi_{r, 2}$ in exactly the same manner as in the two previous sections via the fusion^m

$$\Phi_{h_{r, 0, q}} \otimes \Phi_{h_{1, 2, 0}} \rightarrow \Phi_{h_{r, 2, q}} + \dots \quad (4.14)$$

We obtain for $\Psi_{r, 2}$, $r \in \mathbb{N}$:

$$\mathcal{L}_{-1} = tL_{-1} - tG_{-\frac{1}{2}}^+ G_{-\frac{1}{2}}^- + T_{-1} \quad , \quad \psi_0 = \begin{pmatrix} -t |h_{r, 2, q}\rangle \\ q |h_{r, 2, q}\rangle \\ 0 \\ 0 \end{pmatrix}, \quad (4.15)$$

^mWe still have the freedom to choose $\tilde{q}$ conveniently; we suggest $\tilde{q} = 0$.

$$\mathcal{E}(n) = \begin{pmatrix} -\frac{q}{nt^2(n-r)}\mathcal{L}_{-1} & \frac{1}{nt(n-r)}\mathcal{L}_{-1} & -\frac{t(2n-r-1)-2q}{2nt(n-r)}G_{-\frac{1}{2}}^+ & -\frac{t(2n-r-1)+2q}{2nt(n-r)}G_{-\frac{1}{2}}^- \\ \frac{nt^2(n-r)+q^2}{nt^3(n-r)}\mathcal{L}_{-1} & -\frac{q}{nt^2(n-r)}\mathcal{L}_{-1} & \left(\frac{qt(2n-r-1)-2q^2}{2nt^2(n-r)}-1\right)G_{-\frac{1}{2}}^+ & \left(\frac{qt(2n-r-1)+2q^2}{2nt^2(n-r)}+1\right)G_{-\frac{1}{2}}^- \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix} \quad (4.16)$$

$$\mathcal{T}(s) = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ \frac{-t(r-2s)}{2(q-1)+t(r-2s)}G_{-\frac{1}{2}}^- & \frac{2t}{2(q-1)+t(r-2s)}G_{-\frac{1}{2}}^- & \frac{2}{2(q-1)+t(r-2s)}\mathcal{L}_{-1} & 0 \\ \frac{t(r-2s)}{2(q+1)-t(r-2s)}G_{-\frac{1}{2}}^+ & \frac{2t}{2(q+1)-t(r-2s)}G_{-\frac{1}{2}}^+ & 0 & \frac{2}{2(q+1)-t(r-2s)}\mathcal{L}_{-1} \end{pmatrix} \quad (4.17)$$

$$\mathcal{W}(n) = \left(-\frac{q}{t}\mathcal{L}_{-1}, \mathcal{L}_{-1}, -\left(\frac{t}{2}(n-1)-q\right)G_{-\frac{1}{2}}^+, -\left(\frac{t}{2}(n-1)+q\right)G_{-\frac{1}{2}}^- \right) \quad (4.18)$$

$$\Psi_{r,2} = \mathcal{W}(r)\mathcal{T}(r-\frac{1}{2})\mathcal{E}(r-1)\mathcal{T}(r-\frac{3}{2})\mathcal{E}(r-2)\dots\mathcal{E}(1)\mathcal{T}(\frac{1}{2})\psi_0. \quad (4.19)$$

4.4. General uncharged singular vectors

Based on associativity arguments for the fusions found in the previous sections we conjecture that the fusion

$$\Phi_{h_{0,s+2},q-\tilde{q}} \otimes \Phi_{h_{r+1,2},\tilde{q}} \rightarrow \Phi_{h_{r,s},q} + \dots \quad (4.20)$$

will give us an expression for the uncharged singular vector $\Psi_{r,s}$, where we have to use the knowledge of the singular vector $\Psi_{r,2}$ obtained in section 4.3.. Another possible fusion we could take is

$$\Phi_{h_{r,0},q-\tilde{q}} \otimes \Phi_{h_{1,s},\tilde{q}} \rightarrow \Phi_{h_{r,s},q} + \dots, \quad (4.21)$$

where $\Psi_{1,s}$ can be found using the fusion $\Phi_{h_{0,s+2},q} \otimes \Phi_{h_{2,2},0} \rightarrow \Phi_{h_{1,s},q} + \dots$. However, $\Psi_{2,2}$ is at level two, which makes the recursion procedure much more complicated.

5. Changing the fusion point

The fusion we were using in the last sections corresponds to the three point function taken at the super point $Z = (z, \theta^+, \theta^-)$ and the origin:

$$\langle 0 | \Phi_{h_j,q_j}(0) \Phi_{h_1,q_1}(Z) \Phi_{h_2,q_2}(0) | 0 \rangle. \quad (5.1)$$

This choice was made for simplicity, however we could have chosen a three point function which is based on three different points:

$$\langle 0 | \Phi_{h_j, q_j}(Z_f) \Phi_{h_1, q_1}(Z_1) \Phi_{h_2, q_2}(Z_2) | 0 \rangle \quad . \quad (5.2)$$

In fact choosing the mid point of Z_1 and Z_2 as fusion point Z_f was a crucial step in the Virasoro case to calculate the precise formulae of Benoit and Saint-Aubin¹³. The same trick also works for the affine algebra²³ $A_1^{(1)}$. Although the singular vectors are independent of the chosen fusion point, the recursion matrices depend on it. In this section we investigate this dependence and calculate it explicitly for the vectors $\Psi_{r,2}$ choosing the mid point as fusion point.

5.1. Operator product expansion

Exactly as before we need to find the action of the algebra generators on the coefficients of the operator product expansion, the so called descent equations. For simplicity, we will only choose fusion points Z_f on the line connecting the two super points Z_1 and Z_2 . This simplifies the operator product expansion so that it only depends on the super co-ordinates Z_{12} and $\theta_{12}^\pm$:

$$\begin{aligned} \Phi_{h_1, q_1}(Z_1) \Phi_{h_2, q_2}(Z_2) | 0 \rangle &= \sum_{j \in \mathcal{J}} C_{12}^j Z_{12}^{h_j - h_1 - h_2} \left| \psi_j(Z_{12}, \theta_{12}^\pm, Z_f) \right\rangle , \\ \left| \psi_j(Z_{12}, \theta_{12}^\pm, Z_f) \right\rangle &= \sum_{n \in \mathbb{N}_0} Z_{12}^n |h_j + n, q_j\rangle + \theta_{12}^+ \sum_{r \in \mathbb{N}_0 + \frac{1}{2}} Z_{12}^{r - \frac{1}{2}} |h_j + r, q_j - 1\rangle \\ &\quad + \theta_{12}^- \sum_{r \in \mathbb{N}_0 + \frac{1}{2}} Z_{12}^{r - \frac{1}{2}} |h_j + r, q_j + 1\rangle + \theta_{12}^+ \theta_{12}^- \sum_{n \in \mathbb{N}_0} Z_{12}^{n-1} \overline{|h_j + n, q_j\rangle} . \end{aligned} \quad (5.3)$$

It is worth remarking that the vectors $|h_j + n, q_j\rangle$, $|h_j + r, q_j \pm 1\rangle$ and $\overline{|h_j + n, q_j\rangle}$ still depend on the fusion point. Choosing a different fusion point is the result of conjugating the fields by the generators L_{-1} and $G_{-\frac{1}{2}}^\pm$:

$$\Phi(Z) = e^{zL_{-1}} e^{-\theta^+ G_{-1/2}^-} e^{-\theta^- G_{-1/2}^+} \Phi(0) e^{\theta^- G_{-1/2}^+} e^{\theta^+ G_{-1/2}^-} e^{-zL_{-1}} \quad (5.4)$$

The conjugated commutation relations

$$\begin{aligned} [L_m(Z_0), \Phi(Z_2)] &= \left\{ h(m+1) Z_{20}^m + \frac{1}{2}(m+1) Z_{20}^m (\theta_{20}^+ D_2^- + \theta_{20}^- D_2^+) + Z_{20}^{m+1} \partial_{z_2} \right. \\ &\quad \left. + \frac{q}{2} \theta_{20}^+ \theta_{20}^- Z_{20}^{m-1} m(m+1) \right\} \Phi(Z_2) \\ [G_r^\pm(Z_0), \Phi(Z_2)] &= \left\{ 2h(r + \frac{1}{2}) \theta_{20}^\pm Z_{20}^{r - \frac{1}{2}} - Z_{20}^{r + \frac{1}{2}} D_2^\pm \pm \theta_{20}^+ \theta_{20}^- (r + \frac{1}{2}) Z_{20}^{r - \frac{1}{2}} D_2^\pm \right. \\ &\quad \left. + 2\theta_{20}^\pm Z_{20}^{r + \frac{1}{2}} \partial_{z_2} \pm q \theta_{20}^\pm Z_{20}^{r - \frac{1}{2}} (r + \frac{1}{2}) \right\} \Phi(Z_2) \\ [T_m(Z_0), \Phi(Z_2)] &= \left\{ 2h \theta_{20}^+ \theta_{20}^- m Z_{20}^{m-1} + Z_{20}^m (\theta_{20}^- D_2^+ - \theta_{20}^+ D_2^-) + 2\theta_{20}^+ \theta_{20}^- Z_{20}^m \partial_{z_2} \right. \\ &\quad \left. + q Z_{20}^m \right\} \Phi(Z_2), \end{aligned} \quad (5.5)$$

lead us to the conjugated version of the representation σ :

$$\begin{aligned}
\sigma(L_m(Z_0))\Phi(Z_2) &= \left\{ L_m(Z_0) - h(m+1)Z_{20}^m - Z_{20}^{m+1}\partial_{z_2} - \frac{1}{2}(m+1)Z_{20}^m(\theta_{20}^+D_2^- + \theta_{20}^-D_2^+) \right. \\
&\quad \left. - \frac{q}{2}m(m+1)\theta_{20}^+\theta_{20}^-Z_{20}^{m-1} \right\}\Phi(Z_2) \\
\sigma(G_r^\pm(Z_0))\Phi(Z_2) &= \left\{ G_r^\pm(Z_0) - 2h(r + \frac{1}{2})\theta_{20}^\pm Z_{20}^{r-\frac{1}{2}} \mp (r + \frac{1}{2})\theta_{20}^+\theta_{20}^-Z_{20}^{r-\frac{1}{2}}D_2^\pm \right. \\
&\quad \left. + Z_{20}^{r+\frac{1}{2}}D_2^\pm - 2\theta_{20}^\pm Z_{20}^{r+\frac{1}{2}}\partial_{z_2} \mp q(r + \frac{1}{2})Z_{20}^{r-\frac{1}{2}}\theta_{20}^\pm \right\}\Phi(Z_2) \\
\sigma(T_m(Z_0))\Phi(Z_2) &= \left\{ T_m(Z_0) - 2hm\theta_{20}^+\theta_{20}^-Z_{20}^{m-1} - Z_{20}^m(\theta_{20}^-D_2^+ - \theta_{20}^+D_2^-) \right. \\
&\quad \left. - 2\theta_{20}^+\theta_{20}^-Z_{20}^m\partial_{z_2} - qZ_{20}^m \right\}\Phi(Z_2) .
\end{aligned} \tag{5.6}$$

We classify the fusion point by a parameter η which reflects its position on the super line connecting $Z_1 = (z_1, \theta_1^+, \theta_1^-)$ and $Z_2 = (z_2, \theta_2^+, \theta_2^-)$:

$$Z_f = (z_f, \theta_f^+, \theta_f^-) = (z_2 + \eta(z_1 - z_2), \theta_2^+ + \eta(\theta_1^+ - \theta_2^+), \theta_2^- + \eta(\theta_1^- - \theta_2^-)) . \tag{5.7}$$

A simple calculation shows how to relate the super co-ordinates:

$$\begin{aligned}
Z_{1f} &= (1 - \eta)Z_{12} & \theta_{1f}^\pm &= (1 - \eta)\theta_{12}^\pm \\
Z_{2f} &= -\eta Z_{12} & \theta_{2f}^\pm &= -\eta\theta_{12}^\pm .
\end{aligned} \tag{5.8}$$

Taking $\eta = 0$ corresponds to choosing the point Z_2 as fusion point which is exactly what we did in the previous sections, $\eta = \frac{1}{2}$ and $\eta = 1$ correspond to the mid point and Z_1 respectively. For the mid point the coefficients $|h_j + n, q_j\rangle$, $|h_j + r, q_j \pm 1\rangle$ and $|\overline{h_j + n, q_j}\rangle$ only pick up sign factors according to their level by interchanging (h_1, q_1) and (h_2, q_2) , since $Z_{1f} = -Z_{2f}$ and $\theta_{1f}^\pm = -\theta_{2f}^\pm$. This is why the mid point is distinguished.

5.2. Descent equations

The main difference to the descent equations derived above is that by acting with the generators on the OPE we obtain contributions not only from Φ_1 but also from Φ_2 :

$$A(Z_f)\Phi_1(Z_1)\Phi_2(Z_2) = [A(Z_f), \Phi_1(Z_1)]\Phi_2(Z_2) + \Phi_1(Z_1)[A(Z_f), \Phi_2(Z_2)] , \tag{5.9}$$

where A stands for an algebra generator. For $\eta = 0$ it is easy to see that the second part contributes to L_0 and T_0 only. The descent equations for $\eta \in \mathbb{C}$ are:

$$\begin{aligned}
L_m |h_j + n, q_j\rangle &= \left\{ (m+1)(h_1(1-\eta)^m + h_2(-\eta)^m) + ((1-\eta)^{m+1} - (-\eta)^{m+1}) \right. \\
&\quad \left. (h_j - h_1 - h_2 + n - m) \right\} |h_j + n - m, q_j\rangle \\
&\quad + (1-\eta)\eta((1-\eta)^m - (-\eta)^m)L_{-1} |h_j + n - m - 1, q_j\rangle \\
L_m |h_j + r, q_j \pm 1\rangle &= \left\{ (m+1)(h_1(1-\eta)^m + h_2(-\eta)^m) + ((1-\eta)^{m+1} - (-\eta)^{m+1}) \right. \\
&\quad \left. (h_j - h_1 - h_2 + r - \frac{m}{2}) \right\} |h_j + n - m, q_j \pm 1\rangle \\
&\quad - (1-\eta)\eta\frac{m+1}{2}((1-\eta)^m - (-\eta)^m)G_{-\frac{1}{2}}^\pm |h_j + n - m - \frac{1}{2}, q_j\rangle \\
&\quad + (1-\eta)\eta((1-\eta)^m - (-\eta)^m)L_{-1} |h_j + n - m - 1, q_j \pm 1\rangle \\
L_m \overline{|h_j + n, q_j\rangle} &= \left\{ (m+1)(h_1(1-\eta)^m + h_2(-\eta)^m) + ((1-\eta)^{m+1} - (-\eta)^{m+1}) \right. \\
&\quad \left. (h_j - h_1 - h_2 + n) \right\} \overline{|h_j + n - m, q_j\rangle} \\
&\quad + \frac{m(m+1)}{2}(q_1(1-\eta)^{m+1} + q_2(-\eta)^{m+1}) |h_j + n - m, q_j\rangle \\
&\quad - \frac{m+1}{2}(1-\eta)\eta((1-\eta)^m - (-\eta)^m)G_{-\frac{1}{2}}^+ |h_j + n - m - \frac{1}{2}, q_j - 1\rangle \\
&\quad + \frac{m+1}{2}(1-\eta)\eta((1-\eta)^m - (-\eta)^m)G_{-\frac{1}{2}}^- |h_j + n - m - \frac{1}{2}, q_j + 1\rangle \\
&\quad (1-\eta)\eta((1-\eta)^m - (-\eta)^m)L_{-1} \overline{|h_j + n - m - 1, q_j\rangle} \\
G_s^+ |h_j + n, q_j\rangle &= ((-\eta)^{s+\frac{1}{2}} - (1-\eta)^{s+\frac{1}{2}}) |h_j + n - s, q_j + 1\rangle \\
&\quad + (1-\eta)\eta((1-\eta)^{s-\frac{1}{2}} - (-\eta)^{s-\frac{1}{2}})G_{-\frac{1}{2}}^+ |h_j + n - s - \frac{1}{2}, q_j\rangle \\
G_s^+ |h_j + r, q_j - 1\rangle &= -\left\{ 2(s + \frac{1}{2})(h_1(1-\eta)^{s+\frac{1}{2}} + h_2(-\eta)^{s+\frac{1}{2}} + (s + \frac{1}{2}) \right. \\
&\quad (q_1(1-\eta)^{s+\frac{1}{2}} + q_2(-\eta)^{s+\frac{1}{2}}) + (1-2\eta)((1-\eta)^{s+\frac{1}{2}} - (-\eta)^{s+\frac{1}{2}}) \\
&\quad \left. (h_j - h_1 - h_2 + r - s) \right\} |h_j + r - s, q_j\rangle \\
&\quad - ((1-\eta)^{s+\frac{1}{2}} - (-\eta)^{s+\frac{1}{2}}) \overline{|h_j + r - s, q_j\rangle} \\
&\quad - (1-\eta)\eta((1-\eta)^{s+\frac{1}{2}} - (-\eta)^{s+\frac{1}{2}})L_{-1} |h_j + r - s - 1, q_j\rangle \\
&\quad + (1-\eta)\eta((1-\eta)^{s-\frac{1}{2}} - (-\eta)^{s-\frac{1}{2}})G_{-\frac{1}{2}}^+ |h_j + r - s - \frac{1}{2}, q_j - 1\rangle \\
G_s^+ |h_j + r, q_j + 1\rangle &= (1-\eta)\eta((1-\eta)^{s-\frac{1}{2}} - (-\eta)^{s-\frac{1}{2}})G_{-\frac{1}{2}}^+ |h_j + r - s - \frac{1}{2}, q_j + 1\rangle \\
G_s^+ \overline{|h_j + n, q_j\rangle} &= \left\{ 2(s + \frac{1}{2})(h_1(1-\eta)^{s+\frac{1}{2}} + h_2(-\eta)^{s+\frac{1}{2}}) + (s + \frac{1}{2}) \right. \\
&\quad (q_1(1-\eta)^{s+\frac{1}{2}} + q_2(-\eta)^{s+\frac{1}{2}}) + (1-2\eta)((1-\eta)^{s+\frac{1}{2}} - (-\eta)^{s+\frac{1}{2}}) \\
&\quad \left. (h_j - h_1 - h_2 + n - s - \frac{1}{2}) + (s + \frac{1}{2})((1-\eta)^{s+\frac{3}{2}} - (-\eta)^{s+\frac{3}{2}}) \right\} \\
&\quad |h_j + n - s, q_j + 1\rangle \\
&\quad + (1-\eta)\eta((1-\eta)^{s+\frac{1}{2}} - (-\eta)^{s+\frac{1}{2}})L_{-1} |h_j + n - s - 1, q_j + 1\rangle \\
&\quad - (s + \frac{1}{2})(1-\eta)\eta((1-\eta)^{s+\frac{1}{2}} - (-\eta)^{s+\frac{1}{2}})G_{-\frac{1}{2}}^+ \\
&\quad |h_j + n - s - \frac{1}{2}, q_j\rangle \\
&\quad + (1-\eta)\eta((1-\eta)^{s-\frac{1}{2}} - (-\eta)^{s-\frac{1}{2}})G_{-\frac{1}{2}}^+ \overline{|h_j + n - s - \frac{1}{2}, q_j\rangle}
\end{aligned} \tag{5.10}$$

$$\begin{aligned}
G_s^- |h_j + n, q_j\rangle &= ((-\eta)^{s+\frac{1}{2}} - (1-\eta)^{s+\frac{1}{2}}) |h_j + n - s, q_j - 1\rangle \\
&\quad + (1-\eta)\eta((1-\eta)^{s-\frac{1}{2}} - (-\eta)^{s-\frac{1}{2}}) G_{-\frac{1}{2}}^- \left| h_j + n - s - \frac{1}{2}, q_j \right\rangle \\
G_s^- |h_j + r, q_j + 1\rangle &= -\left\{ 2(s + \frac{1}{2})(h_1(1-\eta)^{s+\frac{1}{2}} + h_2(-\eta)^{s+\frac{1}{2}} - (s + \frac{1}{2})) \right. \\
&\quad (q_1(1-\eta)^{s+\frac{1}{2}} + q_2(-\eta)^{s+\frac{1}{2}}) + (1-2\eta)((1-\eta)^{s+\frac{1}{2}} - (-\eta)^{s+\frac{1}{2}}) \\
&\quad (h_j - h_1 - h_2 + r - s) \Big\} |h_j + r - s, q_j\rangle \\
&\quad + ((1-\eta)^{s+\frac{1}{2}} - (-\eta)^{s+\frac{1}{2}}) \overline{|h_j + r - s, q_j\rangle} \\
&\quad - (1-\eta)\eta((1-\eta)^{s+\frac{1}{2}} - (-\eta)^{s+\frac{1}{2}}) L_{-1} |h_j + r - s - 1, q_j\rangle \\
&\quad + (1-\eta)\eta((1-\eta)^{s-\frac{1}{2}} - (-\eta)^{s-\frac{1}{2}}) G_{-\frac{1}{2}}^- \left| h_j + r - s - \frac{1}{2}, q_j + 1 \right\rangle \\
G_s^- |h_j + r, q_j - 1\rangle &= (1-\eta)\eta((1-\eta)^{s-\frac{1}{2}} - (-\eta)^{s-\frac{1}{2}}) G_{-\frac{1}{2}}^- \left| h_j + r - s - \frac{1}{2}, q_j - 1 \right\rangle \\
G_s^- \overline{|h_j + n, q_j\rangle} &= \left\{ -2(s + \frac{1}{2})(h_1(1-\eta)^{s+\frac{1}{2}} + h_2(-\eta)^{s+\frac{1}{2}}) + (s + \frac{1}{2}) \right. \\
&\quad (q_1(1-\eta)^{s+\frac{1}{2}} + q_2(-\eta)^{s+\frac{1}{2}}) - (1-2\eta)((1-\eta)^{s+\frac{1}{2}} - (-\eta)^{s+\frac{1}{2}}) \\
&\quad (h_j - h_1 - h_2 + n - s - \frac{1}{2}) - (s + \frac{1}{2})((1-\eta)^{s+\frac{3}{2}} - (-\eta)^{s+\frac{3}{2}}) \Big\} \\
&\quad |h_j + n - s, q_j - 1\rangle \\
&\quad - (1-\eta)\eta((1-\eta)^{s+\frac{1}{2}} - (-\eta)^{s+\frac{1}{2}}) L_{-1} |h_j + n - s - 1, q_j - 1\rangle \\
&\quad + (s + \frac{1}{2})(1-\eta)\eta((1-\eta)^{s+\frac{1}{2}} - (-\eta)^{s+\frac{1}{2}}) G_{-\frac{1}{2}}^- \\
&\quad \left| h_j + n - s - \frac{1}{2}, q_j \right\rangle \\
&\quad + (1-\eta)\eta((1-\eta)^{s-\frac{1}{2}} - (-\eta)^{s-\frac{1}{2}}) G_{-\frac{1}{2}}^- \overline{|h_j + n - s - \frac{1}{2}, q_j\rangle} \\
T_m |h_j + n, q_j\rangle &= (q_1(1-\eta)^m + q_2(-\eta)^m) |h_j + n - m, q_j\rangle \\
T_m |h_j + r, q_j \pm 1\rangle &= \left\{ q_1(1-\eta)^m + q_2(-\eta)^m \pm ((1-\eta)^{m+1} - (-\eta)^{m+1}) \right\} \\
&\quad |h_j + r - m, q_j \pm 1\rangle \\
&\quad \mp (1-\eta)\eta((1-\eta)^m - (-\eta)^m) G_{-\frac{1}{2}}^\pm \left| h_j + r - m - \frac{1}{2}, q_j \pm 1 \right\rangle \\
T_m \overline{|h_j + n, q_j\rangle} &= \left\{ 2m(h_1(1-\eta)^{m+1} + h_2(-\eta)^{m+1}) - 2(1-\eta)\eta \right. \\
&\quad ((1-\eta)^m - (-\eta)^m)(h_j - h_1 - h_2 + n - m) \Big\} |h_j + n - m, q_j\rangle \\
&\quad + (q_1(1-\eta)^m + q_2(-\eta)^m) \overline{|h_j + n - m, q_j\rangle} \\
&\quad - (1-\eta)\eta((1-\eta)^m - (-\eta)^m) G_{-\frac{1}{2}}^+ \left| h_j + n - m - \frac{1}{2}, q_j - 1 \right\rangle \\
&\quad - (1-\eta)\eta((1-\eta)^m - (-\eta)^m) G_{-\frac{1}{2}}^- \left| h_j + n - m - \frac{1}{2}, q_j + 1 \right\rangle,
\end{aligned}$$

where we have again put $q_j = q_1 + q_2$, $m \in \mathbb{N}$, $n \in \mathbb{N}_0$, $r, s \in \mathbb{N}_0 + \frac{1}{2}$ and all the generators are taken at the fusion point.

5.3. Expansion of the generators

We are now able to describe the algorithm for this more general three point function. We start off taking an allowed fusion which means we take a non trivial three point function

$$\langle 0 | \Phi_{h_j, q_j}(Z_f) \Phi_{h_1, q_1}(Z_1) \Phi_{h_2, q_2}(Z_2) | 0 \rangle. \quad (5.11)$$

We insert as before a known operator mapping $\Phi_{h_2,q_2} |0\rangle$ on a singular vector:

$$\Psi = \mathcal{N}(Z_2) \Phi_{h_2,q_2}(Z_2) |0\rangle. \quad (5.12)$$

Clearly all the generators in $\mathcal{N}$ have to be taken at the point Z_2 . We use the generalised representation (5.6) to obtain again:

$$\begin{aligned} 0 &= \langle 0 | \Phi_{h_j,q_j}(Z_f) \Phi_{h_1,q_1}(Z_1) \mathcal{N}(Z_2) \Phi_{h_2,q_2}(Z_2) |0\rangle \\ &= \langle 0 | \Phi_{h_j,q_j}(Z_f) \sigma(\mathcal{N}(Z_2)) \Phi_{h_1,q_1}(Z_1) \Phi_{h_2,q_2}(Z_2) |0\rangle. \end{aligned} \quad (5.13)$$

We are facing now the problem that we want to act with the generators in $\sigma(\mathcal{N}(Z_2))$ on the OPE of $\Phi_{h_1,q_1}(Z_1) \Phi_{h_2,q_2}(Z_2)$. For this purpose it is necessary to write the generators in $\sigma(\mathcal{N}(Z_2))$ which are taken at the point Z_2 in terms of generators which are taken at the fusion point Z_f .

Using the $N = 2$ Taylor expansion

$$\begin{aligned} f(Z_1) &= \sum_{n=0}^{\infty} \frac{1}{n!} Z_{12}^n \partial_{z_2}^n \left\{ 1 + \theta_{12}^+ D_2^- + \theta_{12}^- D_2^+ + \theta_{12}^+ \theta_{12}^- D_2^+ D_2^- \right. \\ &\quad \left. - \theta_{12}^+ \theta_{12}^- \frac{\partial}{\partial z_2} \right\} f(Z_2), \end{aligned} \quad (5.14)$$

we can expand the equations (2.3) about the fusion point which leads to:

$$\begin{aligned} L_m(Z_2) &= \sum_{n=0}^{\infty} (\eta Z_{12})^n \left\{ \binom{m+1}{n} L_{m-n}(Z_f) - \eta \frac{m+1}{2} \theta_{12}^+ \binom{m}{n} G_{m-n-\frac{1}{2}}^-(Z_f) \right. \\ &\quad \left. - \eta \frac{m+1}{2} \theta_{12}^- \binom{m}{n} G_{m-n-\frac{1}{2}}^+(Z_f) \right. \\ &\quad \left. + \eta^2 \frac{m(m+1)}{2} \binom{m-1}{n} \theta_{12}^+ \theta_{12}^- T_{m-n-1}(Z_f) \right\} \\ G_r^\pm(Z_f) &= \sum_{n=0}^{\infty} (\eta Z_{12})^n \left\{ \binom{r+\frac{1}{2}}{n} G_{r-n}^\pm(Z_f) + 2\eta \theta_{12}^\pm \binom{r+\frac{1}{2}}{n} L_{r-n-\frac{1}{2}}(Z_f) \right. \\ &\quad \left. \pm \eta \theta_{12}^\pm \left(r + \frac{1}{2}\right) \binom{r-\frac{1}{2}}{n} T_{r-n-\frac{1}{2}}(Z_f) \right. \\ &\quad \left. \mp \eta^2 \theta_{12}^+ \theta_{12}^- \left(r + \frac{1}{2}\right) G_{r-n-1}^\pm(Z_f) \right\} \\ T_m(Z_2) &= \sum_{n=0}^{\infty} (\eta Z_{12})^n \left\{ \binom{m}{n} T_{m-n}(Z_f) + \eta \binom{m}{n} \theta_{12}^+ G_{m-n-\frac{1}{2}}(Z_f) \right. \\ &\quad \left. - \eta \binom{m}{n} \theta_{12}^- G_{m-n-\frac{1}{2}}^+(Z_f) + 2\eta^2 \binom{m}{n} \theta_{12}^+ \theta_{12}^- L_{m-n-1}(Z_f) \right\} \end{aligned} \quad (5.15)$$

We replace the generators in $\sigma(\mathcal{N}(Z_2))$ by the generators taken at the fusion point Z_f according to (5.15) and then we follow the recursion procedure exactly as in the previous sections. The only difference is that wherever we find z or $\theta^\pm$ in the expansions $\mathcal{F}$, $\mathcal{J}$ and $\mathcal{S}$ we have to replace them by Z_{12} or $\theta_{12}^\pm$ respectively. The proof follows by conjugation from the case where $\eta = 0$.

5.4. Example

As mentioned earlier, we expect that the coefficients in $\mathcal{F}$ show certain symmetries under (h_1, q_1) and (h_2, q_2) exchange if we choose $\eta = \frac{1}{2}$. For this case we again would like to give the result for the uncharged singular vectors $\Psi_{r,2}$.

It turns out that the determinant of the recursion matrix $\mathcal{S}$ (3.12) does not depend on the parameter η . This reflects the fact that the fusion rules are independent of the chosen fusion point. We again take the singular vector $\Psi_{1,2}$ (2.19) and use the fusion (4.14) to perform the recursion algorithm. In the Virasoro case we would have to deal at level one only with operators $L_{-1}(Z_2)$. Those expanded according to (5.15) stay the same. We see that the generators $G_{-\frac{1}{2}}^{\pm}(Z_2)$ lead to a finite expansion whilst $T_{-1}(Z_2)$ involves an infinite series of $L_{-n}(Z_f)$ and $G_{-r}^{\pm}(Z_f)$ generators. We will not be able to write the vectors $\Psi_{r,2}$ in a closed expression other than a product of matrices, as it was possible in the Virasoro case. However the expressions appear more suitable to apply the “analytic continuation method” to obtain product expressions for all singular vectors.

To write the result in a compact way, we introduce the four component vectors:

$$F_n^+ = \begin{pmatrix} f_n \\ \bar{f}_n \\ f_{n+\frac{1}{2}}^- \\ f_{n+\frac{1}{2}}^+ \end{pmatrix} \quad F_n^- = \begin{pmatrix} \bar{f}_n \\ f_n \\ f_{n-\frac{1}{2}}^- \\ f_{n-\frac{1}{2}}^+ \end{pmatrix} \quad F_0^- = \begin{pmatrix} -t|h_{r,2}, q\rangle \\ q|h_{r,2}, q\rangle \\ 0 \\ 0 \end{pmatrix}. \quad (5.16)$$

We define the matrices:

$$E_1(n) = \frac{1}{4(n-r)nt} \begin{pmatrix} e_1^{11}(n) & e_1^{12}(n) & e_1^{13}(n) & e_1^{14}(n) \\ e_1^{21}(n) & e_1^{22}(n) & e_1^{23}(n) & e_1^{24}(n) \\ 0 & 0 & 4(n-r)nt & 0 \\ 0 & 0 & 0 & 4(n-r)nt \end{pmatrix} \quad (5.17)$$

$$\begin{aligned} e_1^{11}(n) &= -((r+1)t + q + 2 - 2nt)L_{-1} - 4\frac{q}{t}T_{-1} + qG_{-\frac{1}{2}}^+G_{-\frac{1}{2}}^- \\ e_1^{12}(n) &= 4T_{-1} - tG_{-\frac{1}{2}}^+G_{-\frac{1}{2}}^- + tL_{-1} \\ e_1^{13}(n) &= ((r+1)t + 2(q+1) - 2nt)G_{-\frac{1}{2}}^+ \\ e_1^{14}(n) &= ((r+1)t - 2(q-1) - 2nt)G_{-\frac{1}{2}}^- \\ e_1^{21}(n) &= (-(tr + 2q)n + (r+1)q + (q+2)\frac{q}{t} + n^2t)L_{-1} + (n(n-r) + \frac{q^2}{t^2}) \\ &\quad (4T_{-1} - tG_{-\frac{1}{2}}^+G_{-\frac{1}{2}}^-) \\ e_1^{22}(n) &= -4\frac{q}{t}T_{-1} + qG_{-\frac{1}{2}}^+G_{-\frac{1}{2}}^- - qL_{-1} \\ e_1^{23}(n) &= (2(tr + q)n - (r+1)q - 2(q+1)\frac{q}{t} - 2n^2t)G_{-\frac{1}{2}}^+ \\ e_1^{24}(n) &= (-2(tr - q)n - (r+1)q + 2(q-1)\frac{q}{t} + 2n^2t)G_{-\frac{1}{2}}^- \end{aligned}$$

$$E_2(n) = \frac{1}{2(n-r)nt} \begin{pmatrix} L_{-2} - \frac{t}{8}L_{-1}^2 + \frac{q}{t}T_{-2} & -T_{-2} & -G_{-\frac{3}{2}}^+ + \frac{t}{4}L_{-1}G_{-\frac{1}{2}}^+ & -G_{-\frac{3}{2}}^- + \frac{t}{4}L_{-1}G_{-\frac{1}{2}}^- \\ -\frac{q}{t}L_{-2} + \frac{q}{8}L_{-1}^2 - \frac{n(n-r)t^2+q^2}{t^2}T_{-2} & \frac{q}{t}T_{-2} & \frac{q}{t}G_{-\frac{3}{2}}^+ - \frac{q}{4}L_{-1}G_{-\frac{1}{2}}^+ & \frac{q}{t}G_{-\frac{3}{2}}^- - \frac{q}{4}L_{-1}G_{-\frac{1}{2}}^- \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix} \quad (5.18)$$

$$E_k(n) = \frac{(-1)^k}{2^{k-1}(n-r)nt} \begin{pmatrix} L_{-k} + \frac{q}{t}T_{-k} & -T_{-k} & -G_{-k+\frac{1}{2}}^+ & -G_{-k+\frac{1}{2}}^- \\ -\frac{q}{t}L_{-k} - \frac{n(n-r)t^2+q^2}{t^2}T_{-k} & \frac{q}{t}T_{-k} & \frac{q}{t}G_{-k+\frac{1}{2}}^+ & \frac{q}{t}G_{-k+\frac{1}{2}}^- \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix}, \quad k \geq 3 \quad (5.19)$$

$$T_1(s) = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ -\frac{(2st-tr-2)G_{-\frac{1}{2}}^-}{2(2ts-tr-2q+2)} & -\frac{tG_{-\frac{1}{2}}^-}{2ts-tr-2q+2} & -\frac{4T_{-1}-tG_{-\frac{1}{2}}^+G_{-\frac{1}{2}}^-}{2(2ts-tr-2q+2)} & 0 \\ -\frac{(2st-tr-2)G_{-\frac{1}{2}}^+}{2(2ts-tr+2q+2)} & \frac{tG_{-\frac{1}{2}}^+}{2ts-tr+2q+2} & 0 & \frac{4T_{-1}-tG_{-\frac{1}{2}}^+G_{-\frac{1}{2}}^-+2tL_{-1}}{2(2ts-tr+2q+2)} \end{pmatrix} \quad (5.20)$$

$$T_2(s) = \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ \frac{tL_{-1}G_{-\frac{1}{2}}^- - 4G_{-\frac{3}{2}}^-}{4(2ts-tr-2q+2)} & 0 & \frac{T_{-2}}{2ts-tr-2q+2} & 0 \\ \frac{tL_{-1}G_{-\frac{1}{2}}^+ - 4G_{-\frac{3}{2}}^+}{4(2ts-tr+2q+2)} & 0 & 0 & -\frac{T_{-2}}{2ts-tr+2q+2} \end{pmatrix} \quad (5.21)$$

$$T_k(s) = \left(-\frac{1}{2}\right)^{k-2} \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ -\frac{G_{-k+\frac{1}{2}}^-}{2ts-tr-2q+2} & 0 & \frac{T_{-k}}{2ts-tr-2q+2} & 0 \\ -\frac{G_{-k+\frac{1}{2}}^+}{2ts-tr+2q+2} & 0 & 0 & -\frac{T_{-k}}{2st-tr+2q+2} \end{pmatrix}, \quad k \geq 3 \quad (5.22)$$

$$W_1(r) = \begin{pmatrix} -\frac{t}{2}((1-r)t+q+2)L_{-1} - 2qT_{-1} + \frac{t}{2}qG_{-\frac{1}{2}}^+G_{-\frac{1}{2}}^- \\ 2tT_{-1} - \frac{t^2}{2}G_{-\frac{1}{2}}^+G_{-\frac{1}{2}}^- + \frac{t^2}{2}L_{-1} \\ \frac{t}{2}((1-r)t+2(q+1))G_{-\frac{1}{2}}^+ \\ \frac{t}{2}((1-r)t-2(q-1))G_{-\frac{1}{2}}^- \end{pmatrix} \quad (5.23)$$

$$W_2(r) = \begin{pmatrix} -\frac{t^2}{8}L_{-1}^2 + tL_{-2} + qT_{-2} \\ -tT_{-2} \\ -tG_{-\frac{3}{2}}^+ + \frac{t^2}{4}L_{-1}G_{-\frac{1}{2}}^+ \\ -tG_{-\frac{3}{2}}^- + \frac{t^2}{4}L_{-1}G_{-\frac{1}{2}}^- \end{pmatrix} \quad (5.24)$$

$$W_k(r) = \left(-\frac{1}{2}\right)^{k-2} \begin{pmatrix} tL_{-k} + qT_{-k} \\ -tT_{-k} \\ -tG_{-k+\frac{1}{2}}^+ \\ -tG_{-k+\frac{1}{2}}^- \end{pmatrix}, \quad k \geq 3. \quad (5.25)$$

These matrices enable us to write $F_n^\pm$ as:

$$F_n^- = E_1(n)F_{n-1}^+ + E_2(n-1)F_{n-2}^+ + \dots + E_n(n)F_0^+ \quad (5.26)$$

$$F_{n-1}^+ = T_1(n - \frac{1}{2})F_{n-1}^- + T_2(n - \frac{1}{2})F_{n-2}^- + \dots + T_n(n - \frac{1}{2})F_0^- . \quad (5.27)$$

Hence:

$$\Psi_{r,2} = W_1^T(r)F_{r-1}^+ + W_2^T(r)F_{r-2}^+ + \dots + W_r^T(r)F_0^+ . \quad (5.28)$$

If we introduce the matrices

$$E^{(n)} = \begin{pmatrix} E_1(n) & E_2(n) & \dots & E_n(n) \\ \mathbb{O} & E_1(n-1) & \dots & E_{n-1}(n-1) \\ \vdots & \dots & \dots & \vdots \\ \mathbb{O} & \mathbb{O} & \dots & E_1(1) \end{pmatrix} \quad (5.29)$$

$$T^{(n)} = \begin{pmatrix} T_1(n - \frac{1}{2}) & T_2(n - \frac{1}{2}) & \dots & T_n(n - \frac{1}{2}) \\ \mathbb{O} & T_1(n - \frac{3}{2}) & \dots & T_{n-1}(n - \frac{3}{2}) \\ \vdots & \dots & \dots & \vdots \\ \mathbb{O} & \mathbb{O} & \dots & T_1(\frac{1}{2}) \end{pmatrix} \quad (5.30)$$

$$W^{(r)} = \begin{pmatrix} W_1^T(r) & W_2^T(r) & \dots & W_r^T(r) \end{pmatrix}, \quad (5.31)$$

then we are able to write the recursion as a product of these matrices:

$$\begin{pmatrix} F_n^- \\ F_{n-1}^- \\ \vdots \\ F_1^- \end{pmatrix} = E^{(n)} \begin{pmatrix} F_{n-1}^+ \\ F_{n-2}^+ \\ \vdots \\ F_0^+ \end{pmatrix}, \quad \begin{pmatrix} F_{n-1}^+ \\ F_{n-2}^+ \\ \vdots \\ F_0^+ \end{pmatrix} = T^{(n)} \begin{pmatrix} F_{n-1}^- \\ F_{n-2}^- \\ \vdots \\ F_0^- \end{pmatrix} \quad (5.32)$$

$$\Psi_{r,2} = W^{(r)} \begin{pmatrix} F_{r-1}^+ \\ F_{r-2}^+ \\ \vdots \\ F_0^+ \end{pmatrix}. \quad (5.33)$$

Finally, we can write the singular vector in the formⁿ:

$$\begin{aligned} \Psi_{r,2} &= W^{(r)}T^{(r)} \left\{ 1 + P^{(r)}E^{(r)}T^{(r)} + (P^{(r)}E^{(r)}T^{(r)})^2 + \dots + (P^{(r)}E^{(r)}T^{(r)})^{r-1} \right\} \begin{pmatrix} 0 \\ 0 \\ \vdots \\ 0 \\ F_0^- \end{pmatrix} \\ &= W^{(r)}T^{(r)} (1 - P^{(r)}E^{(r)}T^{(r)})^{-1} \begin{pmatrix} 0 \\ 0 \\ \vdots \\ 0 \\ F_0^- \end{pmatrix}, \end{aligned} \quad (5.34)$$

where $P^{(r)}$ is the r by r matrix

$$P^{(r)} = \begin{pmatrix} \mathbb{O} & \mathbb{1} & \mathbb{O} & \dots & \mathbb{O} \\ \mathbb{O} & \mathbb{O} & \mathbb{1} & \dots & \mathbb{O} \\ \vdots & & & & \vdots \\ \mathbb{O} & \mathbb{O} & \mathbb{O} & \dots & \mathbb{1} \\ \mathbb{O} & \mathbb{O} & \mathbb{O} & \dots & \mathbb{O} \end{pmatrix}. \quad (5.35)$$

6. Conclusions

We have shown that the “fusion method” to derive singular vectors can be applied successfully for the $N = 2$ (untwisted) algebra. We have given explicit expressions for all the charged singular vectors Ψ_k and for some of the uncharged singular vectors $\Psi_{r,2}$, and have conjectured the fusion which can be taken to find every uncharged singular vector $\Psi_{r,s}$. However the construction is getting rather complicated. Alternatively, one can use the “analytic continuation method” in the spirit of ref. 16 to find product formulae for $\Psi_{r,s}$ in terms of $\Psi_{n,2}$ expressions. We were not able to start from a charged singular vector and use a similar procedure. The “fermionic” character of a charged singular vector is responsible for the fact that the equations (3.7) lose terms which are necessary to obtain the recursion formulae.

Acknowledgements

I am very grateful to my PhD supervisor Adrian Kent for many discussions about various aspects of this work. I would like to thank Matthias Gaberdiel and Gérard Watts for explaining $N = 2$ fusion and the $N = 1$ Ramond singular vectors to me. I also acknowledge useful discussions with Zoltan Bajnok about the “fusion method”. Furthermore I am very grateful to the referee for pointing out the possibility of changing the fusion point. Finally, and especially,

ⁿIt is worth remarking that $P^{(r)}E^{(r)}$ does not contain $E_k(r)$.

my thanks go to Valérie Blanchard for proofreading the preprint. This work was financially supported in part by SERC and by Greta & Ludwig Dörrzapf.

References

1. H. Ooguri and C. Vafa, Nucl. Phys. B361 (1991) 469
2. H. Nishino, S.J. Gates Jr. and S.V. Ketov, Phys. Lett. B307 (1993) 331
3. M.F. Atiyah, unpublished;
R.S. Ward, Phil. Trans. Roy. Soc. London A315 (1985) 451
4. H. Nishino, Phys. Lett. B324 (1994) 315
5. H.J. Matschull and H. Nicolai, Nucl. Phys. B411 (1994) 609
6. A.A. Belavin, A.M. Polyakov and A.B. Zamolodchikov Nucl. Phys. B241 (1984) 333
7. D. Friedan, Z. Qiu, S. Shenker, Phys. Rev. Lett. Vol. 52 (1984) 1575
8. D. Huse, Phys. Rev. B30 (1984) 3908
9. G.E. Andrews, R.J. Baxter and P.J. Forrester, J. Stat. Phys. 35 (1984) 193
10. M.J. Tejwari, O. Ferreira and O.E. Vilches, Phys. Rev. Lett. Vol. 44 (1980) 152
11. D. Friedan, Z. Qiu, S. Shenker, Phys. Lett. B151 (1985) 37
12. G. Waterson, Phys. Lett. B171 (1986) 77
13. L. Benoit and Y. Saint-Aubin, Phys. Lett. B215 (1988) 517
14. M. Bauer, P. di Francesco, C. Itzykson and J.-B. Zuber, Phys. Lett. B260 (1991) 323
15. M. Bauer, P. di Francesco, C. Itzykson and J.-B. Zuber, Nucl. Phys. B362 (1991) 515
16. A. Kent, Phys. Lett. B273 (1991) 56
17. F.G. Malikov, B.L. Feigin and D.B. Fuchs, Funct. Anal. Appl. Vol. 20 (1986) 103
18. A.Ch. Ganchev and V.B. Petkova, Phys. Lett. B293 (1992) 56;
A.Ch. Ganchev and V.B. Petkova, Phys. Lett. B318 (1993) 77
19. L. Benoit and Y. Saint-Aubin, Int. J. of Mod. Phys. A7 (1992) 3023;
L. Benoit and Y. Saint-Aubin, Lett. Math. Phys. Vol. 23 (1991) 117
20. C.-S. Huang, D.-H. Zhang and Q.-R. Zheng, Nucl. Phys. B389 (1993) 81
21. G.M.T. Watts, Nucl.Phys. B407 (1993) 213
22. P. Bowcock and G.M.T. Watts, Phys. Lett. B297 (1992) 282

- 23. M. Bauer, N. Sochen, Phys. Lett. B275 (1992) 82;
M. Bauer, N. Sochen, Comm. Math. Phys. 152 (1993) 127
- 24. Z. Bajnok, Phys. Lett. B320 (1994) 36
- 25. Z. Bajnok, Phys. Lett. B329 (1994) 225
- 26. J.D. Cohn, Nucl. Phys. B284 (1987) 349
- 27. A. Schwimmer and N. Seiberg, Phys. Lett. B184 (1987) 191
- 28. A.M. Semikhatov, preprint, hep-th/9311180
- 29. E.B. Kiritsis, Phys. Rev. D36 (1987) 3048
- 30. G. Mussardo, G. Sotkov, M. Stanishkov, Int. Journ. of Mod. Phys. A4 (1989) 1135
- 31. M.R. Gaberdiel, Nucl. Phys. B417 (1994) 130
- 32. W. Boucher, D. Friedan, A. Kent, Phys. Lett. B172 (1986) 316